



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04025035.9

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr.:
Application no.: 04025035.9
Demande no:

Anmeldetag:
Date of filing: 21.10.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

B&B Beheer NV
Alkerstraat 28
3570 Alken
BELGIQUE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Use of D4 and 5-HT2A antagonists, inverse agonists or partial agonists

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

EP/02.12.03/EP 03447279
EP/05.01.04/EP 04447001
EP/18.03.04/EP 04447066

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K31/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PL PT RO SE SI SK TR LI

USE OF D4 AND 5-HT2A ANTAGONISTS, INVERSE AGONISTS OR PARTIAL AGONISTS

Field of the invention

The invention relates to the field of neuropsychiatry. More specifically, the invention relates to the use of compounds, which have D4 and 5-HT2A antagonist, inverse agonist or partial agonist activity, for the preparation of medicaments.

Background of the invention

Conventionally, mental disorders are divided into types based on criteria sets with defining features. DSM-IV (*American Psychiatric Association*, (1993 – ISBN 0 – 89042 – 061 – 0)) is the in the art well-known golden standard of such a categorical classification. In DSM-IV, there is no assumption that each category of mental disorder is a completely discrete entity with absolute boundaries dividing it from other mental disorders or from no mental disorder. There is also no assumption that all individuals described as having the same mental disorder are alike in all important ways. Individuals sharing a diagnosis are likely to be heterogeneous even in regard to the defining features of the diagnosis. Thus, the categorical defined mental disorders as mood and anxiety disorders are having an external and even internal variable co-incidence of symptoms concerning e.g. mood, anxiety, perception, feeding, somatic sensations, sexual functions, sleep, cognitive functioning, impulse control, attention, substance use, personality, bereavement, identity, phase of life, abuse or neglect and other aspects of behavior.

In a dimensional system, clinical presentations are classified based on quantification of attributes i.e. dysfunctions rather than the assignment to categories and works best in describing phenomena that are distributed continuously and that do not have clear boundaries.

Emotion dysregulation is known as such an attribution or dysfunction that plays an important role in the development and course of mental disorders (*Gross, J. J. & Munoz, R. F., 1995, Emotion regulation and mental health, Clinical Psychology: Science and Practice, 2, 151-164; Mennin, D.S., Heimberg, R. G., Turk, C. L. & Fresco, D. M., 2002, Applying an emotion regulation framework to integrative approaches to generalized anxiety disorder, Clinical Psychology: Science and Practice, 9, 85-90; Linehan, M. M., 1993, Cognitive-behavioral treatment of borderline personality disorder, New York, The Guilford Press; Gratz, K. L., Roemer, L., 2001 & 2004, Multidimensional assessment of emotion regulation and dysregulation: development, factor structure, and initial validation*

of the Difficulties in Emotion Regulation Scale, Annual meeting of the Association for Advancement of Behavior Therapy, Nov. 2001 & Journal of Psychopathology and Behavioral Assessment, Vol. 26, No. 1, March 2004) besides behavioural and cognitive dysfunctions. D4 dopamine receptors (D4DR), almost exclusively present in the mesocortical and mesolimbic systems (O'Malley, K. L., Harmon, S., Tang, L., Todd, R. D.,
5 The rat dopamine D4 receptor: sequence, gene structure, and demonstration of expression in the cardiovascular system, New Biol., 4, 137-46, 1992), are in the art known as modulators of emotion and cognition. D4DR agonistic activity gives a behavioural sensitisation; D4DR antagonistic activity leads to an emotion modulation (Svensson, T. H.,
10 Mathé, A. A., Monoaminergic Transmitter Systems, Biological Psychiatry (eds. D'Haenen, H., et al.), 45-66, 2002, John Wiley & Sons, Ltd). Data demonstrate that agonism of the dopamine D4 receptors play an important role in the induction of behavioral sensitization to amphetamine and accompanying adaptations in pre- and postsynaptic neural systems associated with the mesolimbocortical dopamine projections (D. L.
15 Feldpausch et al.; The Journal of Pharmacology and Experimental Therapeutics Vol. 286, Issue 1, 497-508, July 1998).

Results suggest that the antagonisms of cortical D2 dopamine receptors are a common target of traditional and atypical antipsychotics for therapeutic action. Higher *in vivo*
20 binding to the D2 receptors in the cortex than in the basal ganglia is suggested as an indicator of favorable profile for a putative antipsychotic compound (X. Xiberas and J.L. Martinot; The British Journal of Psychiatry (2001) 179: 503-508). Results show that dopamine D4 receptor antagonism in the brain does not result in the same neurochemical consequences (increased dopamine metabolism or hyperprolactinemia) observed with
... typical neuroleptics (Smita Patel et al., The Journal of Pharmacology and Experimental Therapeutics Vol. 283, Issue 2, 636-647, 1997). The selective D4 dopamine receptor antagonist L-745,870 was ineffective as an antipsychotic for the treatment of neuroleptic responsive patients with acute schizophrenia (Kramer, M.S. et al., Arch. Gen. Psychiatry 1997 Dec; 54(12):1080).

30 Finally, in the biological system, mental disorders are defined on other levels of abstraction than in the categorical and dimensional system. Structural pathology (e.g. amyloid plaques in Alzheimer Disease), etiology (e.g. HIV Dementia) and deviance from a physiological norm (e.g. reduced cerebral blood flow) are often used as indicative biological markers for a mental disorder. The underlying dysregulation of various
35 neurotransmitter systems (glutaminergic, GABAergic, cholinergic, monoaminergic (nor-

adrenergic, dopaminergic, serotonergic), etc.) is the in the art used model for the explanation of the biological determinants of the clinical presentation of mental disturbances. It is known that the Serotonin 2A Receptor (5-HT_{2A} receptor) - which is widespread in the Central Nervous System (CNS) - has a regulating role on the dysregulation of various neuro-transmitter systems. 5-HT_{2A} agonism gives several behavioural disturbances; 5-HT_{2A} antagonism leads to a governance of mood, social behaviour, anxiety, cognitive function, stress, sleep functions, nociception, sexual functions, feeding and other aspects of behaviour (*J.E. Leysen (2004) 5-HT₂ Receptors; Current Drug Targets – CNS & Neurological Disorders, 2004, 3, 11-26.*

Dysregulation of the HPA axis (hypothalamic - pituitary - adrenal axis) has frequently been reported in patients with psychiatric disorders, and is among the most robustly demonstrated neurobiological changes among psychiatric patients (*D.A. Gutman and C.B. Nemeroff, Neuroendocrinology, Biological Psychiatry (eds. D'Haenen, H., et al), 99, 2002, John Wiley & Sons, Ltd*). The resulting elevated plasma cortisol concentrations leads to an enhanced binding of serotonin for the 5-HT_{2A} receptor (*E. A. Young, Mineralocorticoid Receptor Function in Major Depression, Arch Gen Psychiatry, Jan 2003; 60: 24 - 28*) and thus agonism.

Additionally 5-HT_{2A} antagonism gives a des-inhibiting of the inhibitory effect of the 5-HT_{2A} receptor on (i) the 5-HT_{1A} receptor stimulation by serotonin (*S. M. Stahl, Newer Antidepressants and Mood Stabilizers, Essential Psychopharmacology, 265, University Press; 2 edition (June 15, 2000); ISBN: 0521646154*) and on (ii) the dopamine release in the mesocortical systems (*S. M. Stahl, Classical Antidepressants, Serotonin Selective and Noradrenargic Reuptake Inhibitors, Essential Psychopharmacology, 233, University Press; 2 edition (June 15, 2000); ISBN: 0521646154*).

Clinical or real effectiveness of psychopharma is very rare via common pooping-out; many treatment-refractory patients and up to half of patients fail to attain remission (*S. M. Stahl, Essential Psychopharmacology, Depression and Bipolar Disorders, 151, University Press; 2 edition (June 15, 2000); ISBN: 0521646154*). Implications of not attaining remission for Mental Disorders are increased relapse rates, continuing functional impairment and increased suicide rate (*S. M. Stahl, Essential Psychopharmacology, Depression and Bipolar Disorders, 152, University Press; 2 edition (June 15, 2000); ISBN: 0521646154*). Clinical causes of not attaining remission by the Current Psychopharmacological Compounds are inadequate early treatment, underlying emotion dysregulation (affecting instability – hypersensitivity – hyperaesthesia – dissociative phenomena, etc.) and

competitive antagonism. There is thus a growing need for a more efficient therapy and more efficient, selective and efficacious medicaments for treating mental disorders.

Summary of the invention

- 5 The present invention relates to the use of compounds and pharmaceutical compositions having D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic activity for the treatment of the underlying emotion dysregulation of mental disorders (e.g. affecting instability – hypersensitivity – hyperaesthesia – dissociative phenomena - etc.) and to methods entailing administering to a patient diagnosed as having a mental disorder a
- 10 pharmaceutical composition containing (i) compounds having specific high selective D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic activity and (ii) a known medicinal compound and/or compositions of compounds. The combined D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic effects may reside within the same chemical or biological compound.
- 15 Taken into account the above mentioned (i) rare clinical or real effectiveness of psychotropics, (ii) the governance of the features and dysfunctions responsible - in a variable co-incidentally – for the clinical state of the mental disorders by D4 dopamine receptor (D4DR) and 2A serotonin receptor (5-HT2A) antagonism and (iii) the fact that 5-HT2A antagonism gives a des-inhibiting of the inhibitory effect of the 5-HT2A receptor on (a) the
- 20 5-HT1A receptor stimulation by serotonin and on (b) the dopamine release in the mesocortical systems, the present invention relates to the use of a compound for the preparation of a medicament for treating a disease or disorder with an underlying emotion dysregulation, characterised in that said compound has (i) a selective affinity for the Dopamine-4 (D4) receptor with a pKi value equal to or higher than 8 towards the D4
- 25 receptor and less than 8 towards other Dopamine receptors, and (ii) a selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5-HT receptors and wherein said compound is administered to a patient in a dose ranging between 5 and 15 mg of the active ingredient. Preferably, said compound is pipamperon.
- 30 In a preferred embodiment, in a mono therapeutic context, the invention relates to the use of a compound as defined above, preferably pipamperon, for preparing a medicament for treating a disease or disorder selected from the group comprising anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders, factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment

disorders, cognitive disorders, impulse control disorders, pervasive development, attention-deficit and disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational, identity, phase of life, academic problem, problems related to abuse or neglect.

According to a further embodiment the invention relates to the use of a first compound as defined above for the preparation of a medicament for treating a mental disease or disorder with an underlying emotion dysregulation whereby a second compound is administered simultaneously with, separate from or sequential to said first compound to augment the therapeutic effect of said second compound on said disease, or to provide a faster onset of the therapeutic effect of said second compound on said disease.

The mental diseases or disorders characterized by an underlying emotion dysregulation can be grouped into subclasses as follows: (i) non-cognitive mental disorders comprising mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problems, identity problem, phase of life problem, academic problem and problems related to abuse or neglect; (ii) cognitive diseases comprising delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders; (iii) pain disorders; and (iv) Parkinson Disease.

In a preferred embodiment, the first compound is administered daily at least one day before administering said second compound.

Preferably, said second compound is characterized by the physiological property of influencing positively the activity of the Central Nervous System.

The invention also relates to a method for preparing a compound having a selective D4 and 5-HT2A antagonist, reverse agonist or partial agonist activity comprising the following steps: (a) measuring the selective affinity of a test compound to the D4 receptor and

selecting a compound that has a pKi value equal to or greater than 8 towards the D4 receptor in respect to all the other D receptors, and measuring the selective efficacy of the selected compound to the D4 receptor and selecting a compound which is a selective antagonist, inverse agonist or partial agonist of the D4 receptor; (b) measuring the selective affinity of a test compound to the 5-HT2A receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the 5-HT2A receptor in respect to all the other 5HT receptors, and measuring the selective efficacy of the selected compound to the 5-HT2A receptor and selecting a compound which is a selective antagonist, inverse agonist or partial agonist of the 5-HT2A receptor; (c) identifying a compound which is selected in (a) and (b), (d) preparing the compound identified in (c).

The invention further also relates to a compound prepared by the described method.

Detailed description of the invention

The present inventors surprisingly found that compounds which have a high selective affinity towards the 5-HT2A receptor and which, at the same time have a high selective affinity towards the dopamine-4 (D4) receptor show an improved effect in treating underlying emotion dysregulation of mental disorders.

The compounds according to the invention may be chemical or biological in nature, or may be chemically synthesised. Preferably, the compounds of the invention are provided as a pharmaceutically acceptable salt.

One example of such a compound which has both a selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5HT receptors, and a selective affinity for the D4 receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors is pipamperon. Pipamperon is the conventional name given for the compound of the formula 1'-[3-(p-Fluorobenzoyl)propyl]-[1,4'-bipiperidine]-4'-carboxamide. Pipamperon is also the active ingredient of for instance the commercially available Dipiperon (Janssen, Cilag B.V).

Further, the present inventors surprisingly found that the dosage of active ingredient for pipamperon in treatment (in monotherapy as well as in combination therapy as described in more detail further) could be very low compared to conventionally used dosages. Preferred dosages which, according to the invention, have been shown to be effective for treating these mental disorders, range between 5 and 15 mg per day or between 5 and 10

mg per day. More preferably, dosages of 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 mg per day are used in treatment of the diseases of the invention. In conventional pipamperon treatment, the active ingredient is available in tablets of 40 mg per tablet or in solutions of 2 mg per drop. Conventional usage of high doses ranging from 40 to 360 mg is prescribed. For instance, for children up to the age of 14, doses corresponding with 2 to 6 mg per kg body weight are conventionally prescribed. The high selective affinity of pipamperon towards the 5-HT_{2A} receptor and the D₄ receptor is reflected in the low dosage which is needed for the treatment of the mental diseases listed below and also contributes to the efficacy of the treatment.

- 10 The mental disorders which can be treated using pipamperon in a mono therapy at such low doses are for instance anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders, factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, cognitive disorders, impulse control disorders, pervasive development, attention-deficit and disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational, identity, phase of life, academic problem, problems related to abuse or neglect.

- 20 Mental disorders such as depression are commonly treated with serotonin re-uptake inhibitors. Unfortunately, however, these compounds can give rise to side effects in use. Moreover, a substantial problem in most treatment of mental disorders is the non-response to selective serotonin re-uptake inhibitors (SSRIs). Also the onset of the therapeutic effect can be delayed undesirable.

- 25 A problem to be solved by the present invention is thus the provision of a more efficient therapy and efficient, highly selective and efficacious medicaments for treating mental disorders.

- 30 The inventors found that, for instance, the non-response to selective serotonin re-uptake inhibitors (SSRIs) in depression may be declared by (partial) inhibition of the 5-HT_{1A} stimulation via 5-HT_{2A} stimulation. Des-inhibition thereof via 5-HT_{2A} antagonism seems to be an answer to this problem.

The present inventors found that a simultaneous or foregoing treatment with a compound having a high selective 5-HT_{2A} antagonist, inverse agonist or partial agonist activity, could lead to a greater response towards, for instance, SSRIs. However, not all compounds exhibiting 5-HT_{2A} antagonism are useful: competition between 5-HT_{2A} stimulation via

serotonin and 5-HT2A antagonism via the compound could be responsible for the lack of more efficacy of compounds which have both a selective serotonin re-uptake inhibitory and 5-HT2A antagonist profile, such as trazodone and nefazodone.

5 The present inventors further surprisingly found that a simultaneous or foregoing treatment with a compound having a high selective D4 antagonist, inverse agonist or partial agonist activity in combination with a compound having a high selective 5-HT2A antagonist, inverse agonist or partial agonist activity could lead to a greater response towards, for instance, SSRIs.

10 In this invention, the term "antagonist" refers to an interaction between chemicals in which one partially or completely inhibits the effect of the other, in particular agents having high affinity for a given receptor, but which do not activate this receptor.

In this invention, the term "inverse agonist" refers to a ligand which produces an effect opposite to that of the agonist by occupying the same receptor.

15 In this invention, the term "agonist" relates to an agent which both binds to a receptor and has an intrinsic effect.

In this invention, the term "partial agonist" relates to an agent with lower intrinsic activity than a full agonist, and which produces a lower maximum effect.

20 The present inventors found that a compound which binds to the 5-HT2A receptor with a pKi of at least 8 but for which the binding affinity, i.e. pKi, towards other 5HT receptors is less than 8 in combination with a high selective affinity for the D4 receptor, i.e. which bind to the D4 receptor with a pKi of at least 8 but for which the binding affinity, i.e. pKi, towards other dopamine receptors is less than 8 also show such an improved effect in treatment. These effects, i.e. D4 antagonism, inverse agonism or partial agonism and 5-HT2A antagonism, inverse agonism or partial agonism, may reside in the same
25 compound.

The term "other 5HT receptors" as used herein relate to for instance 5-HT1 receptors (e.g. 5-HT1A, 5-HT1B, 5-HT1D, 5-HT1E, 5-HT1F), 5-HT2B, 5-HT2C, 5-HT6 (rat) and 5-HT7 (rat).

30 By the expression "selective affinity for the 5-HT2A receptor" is meant that the receptor has a higher affinity for the 5-HT2A receptor than for other 5-HT receptors.

The expression "selective affinity for the D4 receptor" means that the receptor has a higher affinity for the dopamine D4 receptor than for other dopamine receptors.

The term "other dopamine receptors" are, for instance, D1, D2 and D3 dopamine receptors.

pKi values of test compounds for dopamine receptors as well as 5-HT_{2A} receptors can be measured using commonly known assays.

- 5 Compounds which have a selective affinity for the D₄ receptor preferably have a pKi value equal to or higher than 8 towards the D₄ receptor and less than 8 towards other dopamine receptors.

Preferably, the compounds of the invention which have a selective affinity for the 5-HT_{2A} receptor (or the D₄ receptor), are compounds which have a pKi value equal to or higher
10 than 8 towards the 5-HT_{2A} receptor and the D₄ receptor, and less than 8 towards other 5-HT receptors or dopamine receptors, respectively, as can be measured, for instance by methods known in the art. For instance, the "NIMH Psychoactive Drug Screening Program (PDSP)" K_i database (<http://kiddb.cwru.edu/nimh/5htp.php>), is a unique resource in the public domain which provides information on the abilities of drugs to interact with an
15 expanding number of molecular targets. The PDSP K_i database serves as a data warehouse for published and internally-derived pKi, or affinity, values for a large number of drugs and drug candidates at an expanding number of G-protein coupled receptors, ion channels, transporters and enzymes. The PDSP internet site also provides for commonly used protocols and assays for measuring pKi values of 5-HT and dopamine receptors.

- 20 A preferred example of a compound which has both a selective affinity for the 5-HT_{2A} receptor with a pKi value equal to or higher than 8 towards the 5-HT_{2A} receptor and less than 8 towards other 5-HT receptors, and a selective affinity for the D₄ receptor with a pKi value equal to or higher than 8 towards the D₄ receptor and less than 8 towards other Dopamine receptors and which is therefore useful in a combination therapy is
25 pipamperon.

Table 1 illustrates the selective affinity of for instance pipamperon for the 5-HT_{2A} and for the D₄ receptor. In addition, Table 1 also illustrates the low or absence of affinity of
pipamperon for other receptors such as the adrenergic receptors Alpha 1A, Alpha 2A, Alpha 2B, Alpha 2C, Beta 1, Beta 2, and the histamine receptor H₁. As such, treating
30 patients with pipamperon will provide for less side effects which otherwise result from simultaneous stimulation of other receptors. Therefore, and according to preferred embodiments, useful compounds according to the invention not only have a selective 5-HT_{2A} and/or D₄ affinity but also a low affinity for other receptors such as the adrenergic and histamine receptors.

The low dosage which can be used in pipamperon treatment, as already described earlier, contributes to the high selective affinity of the compound towards the 5-HT_{2A} receptor and the D₄ receptor and therefore also to the efficacy of the treatment.

The mental diseases or disorders characterized by an underlying emotion dysregulation can be grouped into subclasses as follows: (i) the non-cognitive mental disorders comprising mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problems, identity problem, phase of life problem, academic problem and problems related to abuse or neglect; (ii) cognitive diseases comprising delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders; (iii) the pain disorders; and (iv) Parkinson Disease. In Table 5, this classification has been used for summarizing the diseases and disorders relative to known psychotropics. In Table 6, an overview of pharmacological grouping is provided, indicating the pharmacological profile numbering, the pharmacological profile, the main disease or disorder indication(s), the name of the compound, the dose range, and the company producing or selling said compound.

These diseases and their diagnosis are very clearly defined in the "Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)" published by the American Psychiatric Association. This manual sets forth diagnostic criteria, descriptions and other information to guide the classification and diagnosis of mental disorders and is commonly used in the field of neuropsychiatry. It is for instance available on the internet under: <http://www.behavenet.com/capsules/disorders/dsm4tr.htm>.

The expression "non-cognitive diseases or disorders" used in some of the embodiments of the invention comprises the following group of diseases or disorders: mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders,

impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problems, identity problem, phase of life problem, academic problem and problems related to abuse or neglect.

In other embodiments of the invention, the mental diseases or disorders that are characterized by an underlying emotion dysregulation belong to the group of pain disorders. For instance, the combination therapy with pipamperon is especially advantageous for management of acute pain in diseases such as, but not limited to, musculoskeletal diseases, rheumatoid arthritis, osteoarthritis and ankylosing spondylitis. For the classification of pain disorders, reference is also made to the DSM-IV where these disorders are clearly described in the section of somatoform disorders by way of internationally accepted diagnostic criteria.

In other embodiments of the invention, the 5-HT_{2A} receptor and/or Dopamine-4 receptor antagonist, inverse agonist or partial agonist (e.g. pipamperon) is used in treatment of patients having neuro-degenerative diseases or disorders, or related cognitive diseases or disorders. The diseases or disorders of the present invention are characterized by an underlying degeneration of the Central Nervous System (CNS), preferably selected from the group consisting of, but not limited to, neurodegenerative diseases such as Parkinson Disease, and in other embodiments of the invention, selected from the group of (related) cognitive diseases or disorders such as Alzheimer Disease.

For instance, Parkinson Disease, which is a chronic progressive nervous disease chiefly of later life, is linked to decreased dopamine production in the substantia nigra and is marked by tremor and weakness of resting muscles and by a shuffling gait. Dopamine agonists and even levodopa, widely used in Parkinson Disease, gives via a dopamine D₄ receptor stimulation psychiatric manifestations. The induced release of serotonin acts via 5-HT_{2A} stimulation as a "brake" on dopamine release (Young B.K., Camicioli R., Ganzini L., *Neuropsychiatric adverse effects of antiparkinsonian drugs. Characteristics, evaluation and treatment. Drugs Aging. 1997 May;10(5):367-83*). Because of the need of specific D₄ and 5-HT_{2A} antagonism in the treatment of Parkinson Disease with dopamine agonists and even levodopa, it seems reasonable to combine with a compound with a high selective D₄ and 5-HT_{2A} antagonism i.e. having merely no activity towards the other receptors especially the D₂ receptor because of the primary need of the relieve of the excessive burden of remaining dopaminergic neurons. Therefore, the use of the so-called atypical anti-psychotics or serotonin-dopamine antagonists (SDAs) is absolutely contra-

indicated since their high affinity for the D2 receptor. Even the use of serotonin releasing compounds such as SSRIs in the absence of an effective 5-HT_{2A} antagonism are counterproductive towards the Parkinson Disease symptoms although many Parkinson patients are in need for an antidepressant since major depression is a very common and disabling condition in this kind of patients.

The expression "(related) cognitive diseases or disorders" according to the invention comprises, the following group of diseases or disorders: delirium (F05), dementia (such as Alzheimer Disease (F00), vascular dementia (F01), dementia due to other general medical conditions (HIV disease (F02.4), head trauma (F06.8), Parkinson Disease (F02.3), Huntington Disease (F02.2), Pick Disease (F02.0), Creutzfeldt-Jacob Disease (F02.1) and other (F02.8)), substance-induced persisting dementia (F1x.6)), amnesic disorders due to a general medical condition (F06.8) or a substance-induced persisting amnesic disorder (F1x.6), mild cognitive impairment disorder (F06.7) and other cognitive disorders (F04). The above list of diseases is provided by way of example and is not intended to limit the invention.

For instance, Alzheimer Disease is a degenerative brain disease of unknown cause that is the most common form of dementia. Alzheimer Disease usually starts in late middle age or in old age as a memory loss for recent events spreading to memories for more distant events and progresses over the course of five to ten years to a profound intellectual decline characterized by dementia and personal helplessness. The disease is marked histologically by the degeneration of brain neurons especially in the cerebral cortex and by the presence of neurofibrillary tangles and plaques containing beta-amyloid. Because dopamine receptor D4 (DRD4) antagonism can inhibit the behavioral disturbances – merely aggression and confusion – caused by the degeneration of dopamine D2 receptors (Esiri, M.M., *The basis for behavioural disturbances in dementia*, J. Neurol. Neurosurg. Psychiatry, 1996; 61(2):127–130.2) accompanied with Alzheimer disease and 5-HT_{2A} antagonism has an important boosting effect towards the effect of cholinesterase inhibitors such as used in the treatment by facilitating the affected dopamine release in the mesocortical dopamine pathways, a high selective D4/5-HT_{2A}-antagonist would be a more preferable compound to combine with a cholinesterase inhibitor since this avoids the counteracting effect of the in the art used SDAs on the cognitive functioning by its dopamine receptor D2-antagonism.

These diseases and their diagnoses are very clearly defined in the "*International Statistical Classification of Diseases and Related Health Problems, 1989 Revision, Geneva, World Health Organization, 1992 (ICD-10)*". This manual sets forth diagnostic

criteria, descriptions and other information to guide the classification and diagnosis of neurodegenerative disorders and is commonly used in the field of neurology. According to the ICD-10 classification, the cognitive disorders are classified under several classes of disorders, i.e. dispersed under categories F00 to F19 (see above: respective classification
5 between parentheses). Following the DSM classification, however, they are grouped in one class of diseases or disorders.

The terms "treatment", "treating", and the like, as used herein include amelioration or elimination of a developed mental disease or condition once it has been established or alleviation of the characteristic symptoms of such disease or condition. As used herein
10 these terms also encompass, depending on the condition of the patient, preventing the onset of a disease or condition or of symptoms associated with a disease or condition, including reducing the severity of a disease or condition or symptoms associated therewith prior to affliction with said disease or condition. Such prevention or reduction prior to affliction refers to administration of the compound or composition of the invention
15 to a patient that is not at the time of administration afflicted with the disease or condition. "Preventing" also encompasses preventing the recurrence or relapse-prevention of a disease or condition or of symptoms associated therewith, for instance after a period of improvement. It should be clear that mental conditions may be responsible for physical complaints. In this respect, the term "treating" also includes prevention of a physical
20 disease or condition or amelioration or elimination of the developed physical disease or condition once it has been established or alleviation of the characteristic symptoms of such conditions.

As used herein, the term "medicament" also encompasses the terms "drug", "therapeutic", "potion" or other terms which are used in the field of medicine to indicate a preparation
25 with therapeutic or prophylactic effect.

The present inventors not only found that the selective 5-HT_{2A} and D₄ antagonists, inverse agonists or partial agonists have an effect in augmenting the therapeutic effect or in providing a faster onset of the therapeutic effect of a diversity of other pharmaceutical compounds, i.e. also named "second compounds" in the present invention, in the
30 treatment of specific diseases or disorders. A few examples of other pharmaceutical compounds whose effects are augmented or where the onset of the effect is fastened upon simultaneous or fore-going treatment with a selective 5-HT_{2A} and D₄ antagonist, preferably pipamperon in a low dose, are nor-epinephrine re-uptake inhibitors, neuroleptic agents, dopamine antagonists, or compounds used for treating or alleviating
35 musculoskeletal diseases or disorders. A further list of other pharmaceutical compounds

or second compounds useful according to the invention is provided in Table 5. It should be clear, given the general applicable character of the invention, that this list of other pharmaceutical compounds is very brief and that the invention should not be restricted to the ones exemplified herein. It should be clear that in the present invention, pipamperon is never to be seen as a "second compound".

According to the invention, it thus has been found that the compounds having a selective 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity as described above are useful for augmenting the therapeutic effect of a second compound on a disease.

According to another embodiment of the invention, it has also been found that the compounds having a selective 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity as described above are useful for providing a faster onset of the therapeutic effect of a second compound on a disease.

From the above it should be clear that the selective 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist is also named 'the first compound' in the embodiments of the invention.

According to the invention, when the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity reside in separate compounds, the term "composition" may be used. Compositions of the invention comprise a first element having (i) a selective affinity for the D₄ receptor with a pK_i value equal to or higher than 8 towards the D₄ receptor and less than 8 towards other dopamine receptors, and a second element having (ii) a selective affinity for the 5-HT_{2A} receptor with a pK_i value equal to or higher than 8 towards the 5-HT_{2A} receptor and less than 8 towards other 5-HT receptors.

The expression "the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist" is used herein to indicate a single compound having both activities or to indicate the composition comprising the activities in separate elements.

It should be clear that when, in the present invention, a composition of separate elements is used instead of a single compound, this composition of separate elements may be used in combination with another, i.e. a second, compound to augment the therapeutic effect of the other, i.e. the second, compound on the same or another disease.

When the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist or the composition comprising both elements and the second compound are administered simultaneously, the compounds or active ingredients may be present in a single pharmaceutical composition or formulation. Alternatively the compounds or active ingredients are administered in separate pharmaceutical compositions or formulations for simultaneous or

separate use. The invention thus also relates to pharmaceutical compositions comprising pipamperon and a second compound of the invention and to the uses of these pharmaceutical compositions.

When the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist or the composition comprising both elements of the invention are administered prior to the second compound as defined, the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist or the composition comprising both elements is administered at least during 1 day prior to said second compound. Preferably, the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist (e.g. pipamperon) or the composition comprising both elements is administered for at least 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 days prior to the administration of the second compound. Preferably, the 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist (e.g. pipamperon) or the composition comprising both elements is administered for at least 2, 3, 4 or 5 weeks prior to the administration of the second compound, or even for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months prior to the administration of the second compound.

According to a preferred embodiment of the invention, the above described compounds or the composition comprising both elements having a 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity are useful for augmenting the therapeutic effect of citalopram or for providing a faster onset of the therapeutic effect of citalopram.

Citalopram or citalopram hydrobromide is a selective serotonin (5-hydroxytryptamine / 5-HT) re-uptake inhibitor (SSRI) and is the conventional name given for the compound of the formula (RS)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalanecarbonitrile hydro-bromide.

According to an embodiment, a daily dose of active ingredient of SSRI, preferably citalopram, ranges between 10 and 40 mg per day. Preferably, daily doses of active ingredient ranging between 20 and 30 mg per day are administered. More preferably, a daily dose of 10, 15, 20, 25, 30, 35 or 40 mg per day is administered.

According to another preferred embodiment of the invention, the above described compounds or the composition comprising both elements having a 5-HT_{2A} and D₄ antagonist, inverse agonist or partial agonist activity are useful for augmenting the therapeutic effect of citalopram or for providing a faster onset of the therapeutic effect of fluvoxamine.

Fluvoxamine or fluvoxamine maleate (luvox, fevarin) is a selective serotonin (5-HT) reuptake inhibitor (SSRI) belonging to a new chemical series, the 2-aminoethyl oxime

ethers of aralkylketones. It is chemically unrelated to other SSRIs and clomipramine. It is chemically designated as 5-methoxy-4'-(trifluoromethyl) valerophenone (E)-O-(2-aminoethyl) oxime maleate (1:1).

5 According to an embodiment, a daily dose of active ingredient of fluvoxamine maleate ranges between 100 and 300 mg per day. Preferably, daily doses of active ingredient ranging between 150 and 200 mg per day are administered. More preferably, a daily dose of 100, 150, 200, 250 or 300 mg per day is administered.

10 Most of the second compounds herein described are known in the art and may be used in doses according to the supplier's or physician's prescription, or may be used according to specific embodiments described herein.

Also encompassed by the invention are pro-drugs to these second compounds or active metabolites of these compounds. For instance, for risperidone it is known that, among other products, bio transformation in the liver produces 9-hydroxyrisperidone, which is of the same pharmacological activity and intensity as parent risperidone. Therefore, also 9-
15 hydroxyrisperidone, naturally produced or chemically synthesized may be used in the methods and uses according to the invention.

The term "active metabolite" as used herein relates to a therapeutically active compound produced by the metabolism of a parent drug. Drugs administered to treat diseases are usually transformed (metabolized) within the body into a variety of related chemical forms (metabolites), some of which may have therapeutic activity (an active metabolite).
20

The present invention also encompasses the use of these second compounds, administered in the form of a pharmaceutically acceptable salt in admixture with a suitable pharmaceutically acceptable excipient.

To prepare the pharmaceutical compositions, comprising the compounds or the
25 combination of the first and second compound described herein, an effective amount of the active ingredients, in acid or base addition salt form or base form, is combined in admixture with a pharmaceutically acceptable carrier, which can take a wide variety of forms depending on the form of ~~preparation~~ desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, for
30 administration orally, nasal, rectally, percutaneously or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders,

disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included.

The pharmaceutical compounds for treatment are intended for parenteral, topical, oral or local administration and generally comprise a pharmaceutically acceptable carrier and an amount of the active ingredient sufficient to reverse or prevent the bad effects of mental disorders. The carrier may be any of those conventionally used and is limited only by chemico-physical considerations, such as solubility and lack of reactivity with the compound, and by the route of administration.

Examples of pharmaceutically acceptable acid addition salts for use in the present inventive pharmaceutical composition include those derived from mineral acids, such as hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric and sulfuric acids, and organic acids, such as tartaric, acetic, citric, malic, lactic, fumaric, benzoic, glycolic, gluconic, succinic, p-toluenesulphonic acids, and arylsulphonic, for example.

The pharmaceutically acceptable excipients described herein, for example, vehicles, adjuvants, carriers or diluents, are well-known to those who are skilled in the art and are readily available to the public. It is preferred that the pharmaceutically acceptable carrier be one that is chemically inert to the active compounds and one that has no detrimental side effects or toxicity under the conditions of use.

The following formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intramuscular, interperitoneal, rectal, and vaginal administration are merely exemplary and are in no way limiting. Overall, the requirements for effective pharmaceutical carriers for parenteral compositions are well known to those of ordinary skill in the art. See *Pharmaceutics and Pharmacy Practice*, J.B. Lippincott Company, Philadelphia, PA, Banker and Chalmers, eds., pages 238-250, (1982), and *ASHP Handbook on Injectable Drugs*, Toissel, 4th ed., pages 622-630 (1986). Topical formulations, including those that are useful for transdermal drug release, are well-known to those of skill in the art and are suitable in the context of the present invention for application to skin.

Formulations suitable for oral administration require extra considerations considering the nature of the compounds and the possible breakdown thereof if such compounds are administered orally without protecting them from the digestive secretions of the

gastrointestinal tract. Such a formulation can consist of (a) liquid solutions, such as an effective amount of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Capsule forms can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of lactose, sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible excipients. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such excipients as are known in the art.

The compounds of the present invention, alone or in combination with other suitable components, can be made into aerosol formulations to be administered via inhalation. For aerosol administration, the compounds are preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of compounds are 0.01%-20% by weight, preferably 1%-10%. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. The surfactant may constitute 0.1%-20% by weight of the compounds, preferably 0.25-5%. The balance of the compounds is ordinarily propellant. A carrier can also be included as desired, e.g., lecithin for intranasal delivery. These aerosol formulations can be placed into acceptable pressurized propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like. They also may be formulated as pharmaceuticals for non-pressured preparations, such as in a nebulizer or an atomizer. Such spray formulations may be used to spray mucosa.

It will be understood that, apart from daily doses, the compounds can be administered by other schedules. For instance, the present invention also contemplates depot injection, in which a long acting form of the active compound is injected into the body, such as the muscles. From there the active compound slowly enters the rest of the body, so one injection can last from 1 to 4 weeks or even multiple months. Other form of dosage administrations relate to "once-a-week" pills, in which the ingredient is slowly released over a period of a week, and slow-release patches, e.g. a CDS (Continuous Delivery System), or Once-a-Day Transdermal Patches.

According to a further embodiment, the invention also relates to a method for preparing a compound or composition having a selective D4 and 5-HT_{2A} antagonist, reverse agonist or partial agonist. The invention also relates to the compounds prepared by the claimed method, with the proviso that said compound is not an already known compound, such as pipamperon.

It should be clear that the compounds and compositions described herein are useful for treating any patient in need thereof. As used herein the term "patient" is not restricted to humans but also to other mammals, for instance, domestic animals which may also suffer from any form of a mental disease or disorder described herein.

The second compounds of the invention can be further grouped according to their pharmacological profile, which is summarized in Table 6.

The present invention is now described in more detail by the following embodiments. The compounds belonging to different pharmacological profiles can be further grouped according to their action on the same pathway or system as follows.

1: combination therapy with a 5-HT (serotonin) reuptake enhancer

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT (serotonin) reuptake enhancer, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
5 factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said
10 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT (serotonin) reuptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT (serotonin) reuptake enhancer compound, further characterized in that pipamperon
15 is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said
20 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT (serotonin) reuptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT (serotonin) reuptake enhancer compound, further characterized in that pipamperon
25 is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT (serotonin) reuptake enhancer compound is tianeptine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.
30 Preferably, tianeptine is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT (serotonin) reuptake enhancer, preferably tianeptine or a pro-drug or an
35 active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying

emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorder, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

A pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT (serotonin) reuptake enhancer is tianeptine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

2: combination therapy with a 5-HT₁ autoreceptor agonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a 5-HT₁ autoreceptor agonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

25

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₁ autoreceptor agonist compound to augment

the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT1 autoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
10 from or prior to the administration of a 5-HT1 autoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT1 autoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT1 autoreceptor agonist compound is sunepitron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

20

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT1 autoreceptor agonist, preferably sunepitron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of
25 diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorder, antisocial behaviour, bereavement, occupational problem, problems related to
30 abuse or neglect and pain disorders.

3: combination therapy with a 5-HT1A (serotonin 1A receptor) agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy
35 with a 5-HT1A (serotonin 1A receptor) agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders,

premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational
5 problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced
10 persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
15 emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-
20 related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic
25 effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention further also relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
30 emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of
35 said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention further also relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group consisting of Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, 10 dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound 15 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 The present invention further also relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound 25 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT_{1A} (serotonin 1A receptor) agonist compound is chosen from the group consisting of MN-305, zalospirone, xaliproden, VPI-013 (also known as OPC-14523), tandospirone, sarizotan, PRX-00023, metanospirone, lesopitron, gepirone, flesinoxan, EMD 68843, buspirone, bupropion (preferably controlled release formulation) 35 and alnespirone, preferably xaliproden, sarizotan, gepirone, flesinoxan and bupropion (preferably controlled release formulation) or a pro-drug or an active metabolite thereof,

or a pharmaceutically acceptable salt thereof. More preferably, said 5-HT1A (serotonin 1A receptor) agonist is xaliproden and is to be administered in a daily dose ranging between 1 and 2 mg of the active ingredient. Even more preferably, said 5-HT1A (serotonin 1A receptor) agonist is bupropion (controlled release formulation) and is to be administered
5 in a daily dose ranging between 150 and 450 mg of the active ingredient. Even more preferably, said 5-HT1A (serotonin 1A receptor) agonist is gepirone and is to be administered in a daily dose, ranging between 20 and 80 mg of the active ingredient per day.

10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT1A (serotonin 1A receptor) agonist, preferably chosen from the group consisting of MN-305, zalospirone, xaliproden, VPI-013 (also known as OPC-14523), tandospirone, sarizotan, PRX-00023, metanospirone, lesopitron, gepirone, flesinoxan, EMD 68843, buspirone, bupropion (preferably controlled release formulation) and
15 alnespirone, more preferably xaliproden, sarizotan, gepirone, flesinoxan and bupropion (preferably controlled release formulation), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders
20 consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-induced persisting
25 dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson
30 Disease.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active
35 ingredient and wherein said 5-HT1A (serotonin 1A receptor) agonist is xaliproden, preferably provided in a unitary dose of between 1 and 2 mg of the active ingredient.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT1A (serotonin 1A receptor) agonist is bupropion
5 (controlled release formulation), preferably provided in a unitary dose of between 150 and 450 mg of the active ingredient.

The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active
10 ingredient and wherein said 5-HT1A (serotonin 1A receptor) agonist is gepirone, preferably provided in a unitary dose of between 20 and 80 mg of the active ingredient.

4: combination therapy with a 5-HT1A (serotonin 1A receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective
15 affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT1A (serotonin 1A receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders,
20 impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
25 emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related
30 disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT1A antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT1A
35 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT1A antagonist compound is chosen from the group consisting of robalzotan tartrate hydrate and NAD299 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT1A antagonist, preferably chosen from the group consisting of robalzotan tartrate hydrate and NAD299, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

5: combination therapy with a 5-HT1B (serotonin 1B receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT2A and D4-receptor, for instance pipamperon, in a combination therapy with a 5-HT1B (serotonin 1B receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders,

- adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with,
- 5 separate from or prior to the administration of a 5-HT1B antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT1B antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 10 According to a preferred embodiment, the invention relates to the use as described above, wherein said 5-HT1B antagonist compound is chosen from the group consisting of elzasonan, AZD1134 and AR-A2, preferably elzasonan, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.
- 15 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT1B antagonist, preferably chosen from the group consisting of elzasonan, AZD1134 and AR-A2, preferably elzasonan, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation
- 20 of a non-cognitive mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-
- 25 related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect.

6: combination therapy with a 5-HT2B (serotonin 2B receptor) antagonist compound

- The mental disorders which can be treated using compounds having a high selective
- 30 affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT2B (serotonin 2B receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders,
- 35 adjustment disorders, impulse control disorders, substance-related disorder, personality

disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2B} antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{2B} antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT2B antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT2B antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said 5-HT_{2B} antagonist compound is agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, agomelatine is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

35 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT_{2B} antagonist, preferably agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for

simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

- 10 A pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT2B antagonist is agomelatine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

15 7: combination therapy with a 5-HT2C (serotonin 2C receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT2C (serotonin 2C receptor) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

25

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT2C antagonist compound to augment the therapeutic effect or to

35

provide a faster onset of the therapeutic effect of said 5-HT_{2C} antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2C} antagonist compound to augment the
- 10 therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{2C} antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- According to a preferred embodiment, the invention relates to the uses as described
- 15 above, wherein said 5-HT_{2C} antagonist compound is chosen from the group consisting of SB 243213 and agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, agomelatine is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

- 20 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT_{2C} antagonist, preferably chosen from the group consisting of SB 243213 and agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or
- 25 disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement,
- 30 occupational problem, problems related to abuse or neglect and pain disorders.

- The present invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said 5-HT_{2C} antagonist is agomelatine, preferably provided in a
- 35 unitary dose of between 25 and 50 mg of the active ingredient.

8: combination therapy with a 5-HT3 (serotonin 3 receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT3 (serotonin 3 receptor) antagonist compound, are substance-related disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance-related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT3 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT3 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said 5-HT3 antagonist compound is ondansetron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, ondansetron is to be administered in a daily dose ranging between 8 and 32 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT3 antagonist, preferably ondansetron or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of substance-related disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient, and wherein said 5-HT3 antagonist is ondansetron, preferably provided in a unitary dose of between 8 and 32 mg of the active ingredient.

9: combination therapy with a 5-HT6 (serotonin 6 receptor) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT2A and D4 receptor, for instance pipamperon, in a combination therapy with a 5-HT6 (serotonin 6 receptor) antagonist compound, are chosen from the group of

diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT6 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT6 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said 5-HT6 antagonist compound is chosen from the group consisting of SB-271046, 742457 and 271046 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a 5-HT6 antagonist, preferably chosen from the group consisting of SB-271046, 742457 and 271046 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-

Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

5 10: combination therapy with an acetylcholinesterase inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an acetylcholinesterase inhibitor compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, 10 vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

15

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, 20 vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders, characterized in that pipamperon or said 25 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an acetylcholinesterase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said acetylcholinesterase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active 30 ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said acetylcholinesterase inhibitor compound is chosen from the group consisting of tacrine, rivastigmine tartrate, rivastigmine, physostigmine, phenserine tartrate, 35 metrifonate, huperzine A, galantamine (preferably extended release formulation), donepezil, dichlorvos and anseculin hydrochloride, preferably tartrate, or a pro-drug or an

active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, rivastigmine tartrate is to be administered in a daily dose ranging between 3 and 12 mg of the active ingredient. Preferably, phenserine tartrate is to be administered in a daily dose ranging between 20 and 30 mg of the active ingredient. Preferably, galantamine (extended release formulation) is to be administered in a daily dose ranging between 8 and 24 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an acetylcholinesterase inhibitor, preferably chosen from the group consisting of tacrine, rivastigmine tartrate, rivastigmine, physostigmine, phenserine tartrate, metrifonate, huperzine A, galantamine (preferably extended release formulation), donepezil, dichlorvos and anseculin hydrochloride, preferably tartrate, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is rivastigmine tartrate, preferably provided in a unitary dose of between 3 and 12 mg of the active ingredient.

The invention further relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is phenserine tartrate, preferably provided in a unitary dose of between 20 and 30 mg of the active ingredient.

In addition, the invention relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is galantamine (preferably

extended release formulation), preferably provided in a unitary dose of between 8 and 24 mg of the active ingredient.

11: combination therapy with an adenosine A2a receptor antagonist compound

- 5 The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an adenosine A2a receptor antagonist compound, is Parkinson disease.

10 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adenosine A2a receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
15 adenosine A2a receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 According to a preferred embodiment, the invention relates to the use as described above, wherein said adenosine A2a receptor antagonist compound is KW-6002 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, KW-6002 is to be administered in a daily dose ranging between 40 and 80 mg of the active ingredient.

25 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an adenosine A2a receptor antagonist, preferably KW-6002 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson disease.

30

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said acetylcholinesterase inhibitor is KW-6002, preferably provided in a unitary dose of between 40 and 80 mg of the active ingredient.

35

12: combination therapy with an adrenergic transmitter releaser

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an adrenergic transmitter releaser, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

10

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adrenergic transmitter releaser compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adrenergic transmitter releaser compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adrenergic transmitter releaser compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adrenergic transmitter releaser compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

According to a preferred embodiment, the invention relates to the uses as described above, wherein said adrenergic transmitter releaser compound is pipoxazole or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, pipoxazole is to be administered in a daily dose ranging between 30 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an adrenergic transmitter releaser, preferably pipoxazole, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said adrenergic transmitter releaser is pipoxazole, preferably provided in a unitary dose of between 30 and 60 mg of the active ingredient.

13: combination therapy with an alpha 1 adrenoreceptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an alpha 1 adrenoreceptor antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating

disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep
disorders, adjustment disorders, impulse control disorders, personality disorders,
bereavement, occupational problem and problems related to abuse or neglect,
5 characterized in that pipamperon or said pharmaceutically acceptable salt thereof is
administered simultaneously with, separate from or prior to the administration of a alpha 1
adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a
faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound,
further characterized in that pipamperon is to be administered to a patient in a daily dose
10 ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of pain disorders, characterized in that pipamperon or said
15 pharmaceutically acceptable salt thereof is administered simultaneously with, separate
from or prior to the administration of an alpha 1 adrenoreceptor antagonist compound to
augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is
to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
20 ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of Parkinson disease, characterized in that pipamperon or said
25 pharmaceutically acceptable salt thereof is administered simultaneously with, separate
from or prior to the administration of an alpha 1 adrenoreceptor antagonist compound to
augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is
to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
30 ingredient.

According to a preferred embodiment, the invention relates to the uses as described
above, wherein said alpha 1 adrenoreceptor antagonist compound is chosen from the
group consisting of SDZ NVI 085 and flesinoxan or a pro-drug or an active metabolite
35 thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an alpha 1 adrenoreceptor antagonist, preferably chosen from the group consisting of SDZ NVI 085 and flesinoxan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and Parkinson disease.

14: combination therapy with an alpha 2 adrenoreceptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an alpha 2 adrenoreceptor antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a alpha 2 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 2 adrenoreceptor antagonist compound, further characterized in that pipamperon is

to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an alpha 2 adrenoreceptor antagonist compound to
10 augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 2 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described
15 above, wherein said alpha 2 adrenoreceptor antagonist compound is chosen from the group consisting of UK-14304, sunepitron, mirtazepine, idazoxan, fluparoxan, A75200 and (R)-A 75200, preferably sunepitron or idazoxan, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, idazoxan is to be administered in a daily dose ranging between 5 and 40 mg of the active ingredient.

20 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an alpha 2 adrenoreceptor antagonist, preferably chosen from the group consisting of UK-14304, sunepitron, mirtazepine, idazoxan, fluparoxan, A75200 and (R)-A 75200, preferably sunepitron or idazoxan, or a pro-drug or an active metabolite thereof,
25 or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious
30 disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

35 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said alpha 2 adrenoreceptor antagonist is Idazoxan, preferably provided in a unitary dose of between 5 and 40 mg of the active ingredient.

15: combination therapy with an AMPA receptor mediator compound

- 5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate) receptor mediator compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual
10 syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem,
15 problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced
20 persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

- The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
25 emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders,
30 disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an
35 AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further

characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 According to a preferred embodiment, the invention relates to the uses as described above, wherein said AMPA receptor mediator compound is chosen from the group consisting of ampakine ORG 24448/CX-619, ampakine CX-717, ampakine CX-691 and ampakine CX-516, preferably ampakine ORG 24448/CX-619, ampakine CX-717 or ampakine CX-691, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

35

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an AMPA receptor mediator, preferably chosen from the group consisting of ampakine ORG 24448/CX-619, ampakine CX-717, ampakine CX-691 and ampakine CX-516, preferably ampakine ORG 24448/CX-619, ampakine CX-717 or ampakine CX-691, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

16: combination therapy with an amphetamine compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an amphetamine compound, are attention-deficit disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of attention deficit disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an amphetamine compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said amphetamine compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said amphetamine compound is methylphenidate (preferably administered by the transdermal system) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

5

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an amphetamine, preferably methylphenidate (preferably administered by the transdermal system) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of attention deficit disorders.

10

17: combination therapy with an amyloid aggregation-inhibitor compound

15

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an amyloid aggregation-inhibitor compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

20

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an amyloid aggregation-inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said amyloid aggregation-inhibitor compound, further

30

35

characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 According to a preferred embodiment, the invention relates to the use as described above, wherein said amyloid aggregation-inhibitor compound is chosen from the group consisting of APAN and Alzhemed, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, Alzhemed is to be administered in a daily dose of between 200 and 300 mg of the active ingredient.

10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an amyloid aggregation-inhibitor, preferably chosen from the group consisting of APAN and Alzhemed, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer
15 Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-
20 induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient
25 and wherein said amyloid aggregation-inhibitor is Alzhemed, preferably provided in a unitary dose of between 200 and 300 mg of the active ingredient.

18: combination therapy with an androgen receptor modulator compound

30 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an androgen receptor modulator compound, are sexual and gender identity disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
35 emotion dysregulation of sexual and gender identity disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered

simultaneously with, separate from or prior to the administration of an androgen receptor modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said androgen receptor modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said androgen receptor modulator compound is LGD2226 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- 10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an androgen receptor modulator, preferably LGD2226 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of sexual and gender identity disorders.

15

19: combination therapy with an beta 3 adrenoreceptor agonist

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an beta 3 adrenoreceptor agonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

25

- The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of

30

35

a beta 3 adrenoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said beta 3 adrenoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
10 from or prior to the administration of an beta 3 adrenoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said beta 3 adrenoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15

According to a preferred embodiment, the invention relates to the uses as described above, wherein said beta 3 adrenoreceptor agonist compound is SR 58611 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

20

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a beta 3 adrenoreceptor agonist, preferably SR 58611 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of
25 diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and
30 pain disorders.

20: combination therapy with a calcium channel modulator compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy
35 with a calcium channel modulator compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia,

vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a calcium channel modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said calcium channel modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a calcium channel modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said calcium channel modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said calcium channel modulator compound is chosen from the group consisting of safinamide and MEM 1003, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a calcium channel modulator, preferably chosen from the group consisting of safinamide and MEM 1003, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson disease.

21: combination therapy with a cannabiod receptor 1 (CB1) antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a cannabiod receptor 1 (CB1) antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannaboid receptor antagonist compound, further characterized in that pipamperon is to
5 be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
10 emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
15 cannaboid receptor 1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
20 emotion dysregulation of delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
25 cannaboid receptor 1 antagonist compound; further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said cannaboid receptor antagonist compound is SR 141716 or a pro-
30 drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a cannaboid receptor antagonist, preferably SR 141716 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined
35 preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of

diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

22: combination therapy with a cathepsin K inhibitor compound

10 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a cathepsin K inhibitor compound, are pain disorders.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cathepsin K inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cathepsin K inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 According to a preferred embodiment, the invention relates to the use as described above, wherein said cathepsin K inhibitor compound is 462795 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

30 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a cathepsin K inhibitor, preferably 462795 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of pain disorders.

23: combination therapy with a choline uptake enhancer compound

35 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a choline uptake enhancer compound, are chosen from the group of diseases or

disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a choline uptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said choline uptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said choline uptake enhancer compound is MKC-231 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, MKC-231 is to be administered in a daily dose of between 20 and 160 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a choline uptake enhancer, preferably MKC-231 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease,

dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

- 5 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said choline uptake enhancer is MKC-231, preferably provided in a unitary dose of between 20 and 160 mg of the active ingredient.

10 24: combination therapy with a COX-2 inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a COX-2 inhibitor compound, are pain disorders.

- 15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a COX-2 inhibitor compound to augment the
20 therapeutic effect or to provide a faster onset of the therapeutic effect of said COX-2 inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- According to a preferred embodiment, the invention relates to the use as described above,
25 wherein said COX-2 inhibitor compound is chosen from the group consisting of valdecoxib, rofecoxib, parecoxib, etoricoxib, COX 189, celecoxib and ABT-963, preferably parecoxib, etoricoxib or COX 189, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, parecoxib is to be administered in a daily dose of between 20 and 80 mg of the active ingredient. Preferably, etoricoxib is to be
30 administered in a daily dose of between 20 and 120 mg of the active ingredient. Preferably, COX 189 is to be administered in a daily dose of between 100 and 800 mg of the active ingredient.

- The invention also relates to a pharmaceutical composition comprising (a) pipamperon,
35 and (b) a COX-2 inhibitor, preferably chosen from the group consisting of valdecoxib, rofecoxib, parecoxib, etoricoxib, COX 189, celecoxib and ABT-963, preferably parecoxib,

etoricoxib or COX 189, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of pain disorders.

5

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-2 inhibitor is parecoxib, preferably provided in a unitary dose of between 20 and 80 mg of the active ingredient.

10

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-2 inhibitor is etoricoxib, preferably provided in a unitary dose of between 20 and 120 mg of the active ingredient.

15

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-2 inhibitor is COX 189, preferably provided in a unitary dose of between 100 and 800 mg of the active ingredient.

20

25: combination therapy with a COX-inhibiting nitric oxide donator (CINOD) compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a COX-inhibiting nitric oxide donator (CINOD) compound, are pain disorders.

25

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a COX-inhibiting nitric oxide donator (CINOD) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said COX-inhibiting nitric oxide donator (CINOD) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

According to a preferred embodiment, the invention relates to the use as described above, wherein said COX-inhibiting nitric oxide donator (CINOD) compound is chosen from the group consisting of AZD4717 and AZD3582 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, AZD3582 is to be administered
5 in a daily dose ranging between 93.75 and 750 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a COX-inhibiting nitric oxide donator (CINOD), preferably chosen from the group consisting of AZD4717 and AZD3582 or a pro-drug or an active metabolite thereof, or a
10 pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein
15 pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said COX-inhibiting nitric oxide donator (CINOD) is AZD3582, preferably provided in a unitary dose of between 93.75 and 750 mg of the active ingredient.

26: combination therapy with a CRF1 (corticot-releasing factor receptor 1) antagonist

20 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a CRF1 (Corticotropin-Releasing Factor receptor 1) antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
25 factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
35 factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders,

personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a CRF1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said CRF1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a CRF1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said CRF1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said CRF1 antagonist compound is chosen from the group consisting of R121919, NBI-34041, elzasonan, CP-448,187, CP-154-526, AAG 561 and 723620, preferably R121919, elzasonan or AAG 561, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, R121919 is to be administered in a daily dose of between 5 and 80 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a CRF1 antagonist, preferably chosen from the group consisting of R121919, NBI-34041, elzasonan, CP-448,187, CP-154-526, AAG 561 and 723620, preferably R121919, elzasonan or AAG 561, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said CRF1 antagonist is R121919, preferably provided in a unitary dose of
5 between 5 and 80 mg of the active ingredient.

27: combination therapy with a D1 (dopamine 1) receptor agonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy
10 with a D1 (dopamine 1) receptor agonist, are chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
15 emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D1 receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
20 D1 receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
25 emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D1 receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D1 receptor agonist compound, further characterized in that pipamperon is to be administered to a
30 patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said D1 receptor agonist compound is DAS-431 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D1 receptor agonist, preferably DAS-431 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

28: combination therapy with D2 (dopamine 2) receptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with D2 (dopamine 2) receptor antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D2 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D2 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
5 from or prior to the administration of a D2 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D2 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
15 from or prior to the administration of a D2 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D2 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described
20 above, wherein said D2 receptor antagonist compound is chosen from the group consisting of bifeprunox, amisulpride aminosultopride and amisulpride, preferably bifeprunox, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

25 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D2 receptor antagonist, preferably chosen from the group consisting of bifeprunox, amisulpride aminosultopride and amisulpride, preferably bifeprunox, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the
30 underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders,
35 personality disorders, psychological factors affecting medical conditions, malingering,

antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

29: combination therapy with D3 (dopamine 3) receptor antagonist

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with D₃ (dopamine 3) receptor antagonist, are chosen from the group of diseases or disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders, delirium and Parkinson disease.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D₃ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D₃ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D₃ receptor antagonist compound to augment the

therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist compound to augment the
- 10 therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor
- 20 agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- According to a preferred embodiment, the invention relates to the uses as described above, wherein said D3 receptor antagonist compound is chosen from the group
- 25 consisting of BSF-201640 and PD 58491, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a D3 receptor antagonist, preferably chosen from the group consisting of BSF-
- 30 201640 and PD 58491, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders,
- 35 dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related

disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders, delirium and Parkinson disease.

5 30: combination therapy with a DA (dopamine) uptake inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a DA (dopamine) uptake inhibitor, are chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

10

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with,
15 separate from or prior to the administration of a DA uptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said DA uptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a DA uptake inhibitor compound to augment the
25 therapeutic effect or to provide a faster onset of the therapeutic effect of said DA uptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, ~~the invention~~ relates to the uses as described
30 above, wherein said DA uptake inhibitor compound is chosen from the group consisting of safinamide and GBR 12909, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon,
35 and (b) a D₂ receptor antagonist, preferably chosen from the group consisting of safinamide and GBR 12909, or a pro-drug or an active metabolite thereof, or a

pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

5

31: combination therapy with an dopamine (receptor) agonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an dopamine (receptor) agonist, are chosen from the group of diseases or disorders
10 consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorders and Parkinson disease.

15

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, eating
20 disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
25 from or prior to the administration of a dopamine (receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine (receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
35 from or prior to the administration of a dopamine (receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said

dopamine (receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine (receptor) agonist compound to augment
- 10 the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine (receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 15 According to a preferred embodiment, the invention relates to the uses as described above, wherein said dopamine (receptor) agonist compound is chosen from the group consisting of sumanirole, SLV 308, sarizotan, S32504, rotigotine (preferably a Once-a-Day Transdermal Patch), ropinirole HCL (preferably controlled-release formulation), pramipexole, DAB452, cabergoline, bromocriptine, alaptide amantadine, bromocriptine,
- 20 cabergoline lisuride and pergolide, preferably sumanirole, rotigotine (preferably a Once-a-Day Transdermal Patch), pergolide or ropinirole HCL (preferably controlled-release formulation), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, sumanirole is to be administered in a daily dose of between 4 and 16 mg of the active ingredient. Preferably, rotigotine (Once-a-Day
- 25 Transdermal Patch) is to be administered in a daily dose of between 4.5 and 13.5 mg of the active ingredient. Preferably, ropinirole HCL (controlled-release formulation) is to be administered in a daily dose of between 0.75 and 24 mg of the active ingredient. Preferably, pergolide is to be administered in a daily dose of between 0.5 and 10 mg of the active ingredient.
- 30 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a dopamine (receptor) agonist, preferably chosen from the group consisting of sumanirole, SLV 308, sarizotan, S32504, rotigotine (preferably a Once-a-Day Transdermal Patch), ropinirole HCL (preferably controlled-release formulation),
- 35 pramipexole, DAB452, cabergoline, bromocriptine, alaptide amantadine, bromocriptine, cabergoline lisuride and pergolide, more preferably sumanirole, rotigotine (preferably a

Once-a-Day Transdermal Patch), ropinirole HCL (preferably controlled-release formulation) or pergolide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorders and Parkinson disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is sumanirole, preferably provided in a unitary dose of between 4 and 16 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is rotigotine (Once-a-Day Transdermal Patch), preferably provided in a unitary dose of between 4.5 and 13.5 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is ropinirole HCL (controlled-release formulation), preferably provided in a unitary dose of between 0.75 and 24 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said dopamine (receptor) agonist is pergolide, preferably provided in a unitary dose of between 0.5 and 10 mg of the active ingredient.

32: combination therapy with a compound activating ERK (extracellular signal-related kinase)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound that activates ERK (extracellular signal-related kinase), are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound that activates ERK (extracellular signal-related kinase) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound that activates ERK (extracellular signal-related kinase), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said compound that activates ERK (extracellular signal-related kinase) is CPI-1189 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, CPI-1189 is to be administered in a daily dose of between 50 and 100 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound that activates ERK (extracellular signal-related kinase), preferably CPI-1189 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said compound that activates ERK (extracellular signal-related kinase) is CPI-1189, preferably provided in a unitary dose of between 50 and 100 mg of the active ingredient.

33: combination therapy with a GABA (gamma-aminobutyric acid) agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a GABA (gamma-aminobutyric acid) agonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders...

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease,

dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA agonist compound is nefiracetam or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GABA agonist, preferably nefiracetam or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

34: combination therapy with a GABA-A agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a GABA-A (gamma-aminobutyric acid receptor A) agonist compound, are sleep disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of sleep disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate

from or prior to the administration of a GABA-A agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA-A agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA-A agonist compound is gaboxadol or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, gaboxadol is to be administered in a daily dose of between 5 and 20 mg of the active ingredient.

10

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an GABA-A agonist, preferably gaboxadol or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of sleep disorders.

15

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A agonist is Gaboxadol, preferably provided in a unitary dose of between 5 and 20 mg of the active ingredient.

20

35: combination therapy with a GABA-A modulator compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a GABA-A (gamma-aminobutyric acid receptor A) modulator compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

25

30

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious

35

disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered
5 simultaneously with, separate from or prior to the administration of a GABA-A modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA-A modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA-A modulator compound is chosen from the group consisting of zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), SL 65.1498, SEP174559, pagoclone, NGD 96-3, indiplon,
15 eszopiclone, CP-730,330 (NGD 96-3) and ocinaplon, preferably zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), pagoclone, NGD 96-3, indiplon or eszopiclone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, zolpidem MR sustained-release version is to be administered in a daily dose of between 10 and 20 mg of the
20 active ingredient. Preferably, zaleplon extended-release is to be administered in a daily dose ranging between 2.5 and 20 mg of the active ingredient. Preferably, pagoclone is to be administered in a daily dose ranging between 7.5 and 60 mg of the active ingredient. Preferably, indiplon is to be administered in a daily dose of between 10 and 20 mg of the active ingredient. Preferably, eszopiclone is to be administered in a daily dose of between
25 2 and 3 mg of the active ingredient. Preferably, ocinaplon is to be administered in a daily dose of between 10 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GABA-A modulator, preferably chosen from the group consisting of zolpidem
30 (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), SL 65.1498, SEP174559, pagoclone, NGD 96-3, indiplon, eszopiclone, CP-730,330 (NGD 96-3) and ocinaplon, preferably zolpidem (preferably MR sustained-release version), zaleplon (preferably extended-release formulation), pagoclone, NGD 96-3, indiplon or eszopiclone, or a pro-drug or an active metabolite thereof, or a
35 pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental

disease or disorder which is chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is zolpidem MR sustained-release version, preferably provided in a unitary dose of between 10 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is zaleplon extended-release, preferably provided in a unitary dose of between 2.5 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is Pagoclone, preferably provided in a unitary dose of between 7.5 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is indiplon, preferably provided in a unitary dose of between 10 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is eszopiclone, preferably provided in a unitary dose of between 2 and 3 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GABA-A modulator is ocinaplon, preferably provided in a unitary dose of between 10 and 60 mg of the active ingredient.

36: combination therapy with a GABA-B antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a GABA-B (gamma-aminobutyric acid receptor B) antagonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA-B antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA-B antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said GABA-B antagonist compound is AVE 7398 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

30

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GABA-B antagonist, preferably AVE 7398 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform

35

disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

5

37: combination therapy with a Glial-cell Line Derived Neurotrophic Factor compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a Glial-cell Line Derived Neurotrophic Factor compound, is Parkinson disease.

10

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a Glial-cell Line Derived Neurotrophic Factor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said Glial-cell Line Derived Neurotrophic Factor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20

According to a preferred embodiment, the invention relates to the use as described above, wherein said Glial-cell Line Derived Neurotrophic Factor compound is GDNF or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, GDNF is to be administered in a daily dose ranging between 3.75 and 30 mg of the active ingredient.

25

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a Glial-cell Line Derived Neurotrophic Factor, preferably GDNF or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson disease.

30

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said Glial-cell Line Derived Neurotrophic Factor is GDNF, preferably provided in a unitary dose of between 3.75 and 30 mg of the active ingredient.

35

38: combination therapy with a glucocorticoid synthesis inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a glucocorticoid synthesis inhibitor compound, are chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glucocorticoid synthesis inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glucocorticoid synthesis inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glucocorticoid synthesis inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glucocorticoid synthesis inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said glucocorticoid synthesis inhibitor compound is metyrapone or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a glucocorticoid synthesis inhibitor, preferably metyrapone or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying

emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of substance related disorders and Parkinson disease.

39: combination therapy with a glutamate receptor antagonist compound

- 5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a glutamate receptor antagonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

- 15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, and problems related to abuse or neglect,
- 20 characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glutamate receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glutamate receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose
- 25 ranging between 5 and 15 mg of the active ingredient.

- The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said
- 30 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a glutamate receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said glutamate receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
- 35 ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said glutamate receptor antagonist compound is LY354740 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- 5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a glutamate receptor antagonist, preferably LY354740 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of
- 10 diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15 40: combination therapy with an GPCR (G-protein-coupled receptor) modulator

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an GPCR (G-protein-coupled receptor) modulator, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders,
- 20 premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

25

- The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating
- 30 disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt
- 35 thereof is administered simultaneously with, separate from or prior to the administration of a GPCR modulator compound to augment the therapeutic effect or to provide a faster

onset of the therapeutic effect of said GPCR modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GPCR modulator compound to augment the
- 10 therapeutic effect or to provide a faster onset of the therapeutic effect of said GPCR modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- According to a preferred embodiment, the invention relates to the uses as described
- 15 above, wherein said GPCR modulator compound is R1204 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GPCR modulator, preferably R1204 or a pro-drug or an active metabolite
- 20 thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative
- 25 disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

30 41: combination therapy with an GR (glucocorticoid receptor) antagonist

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an GR (glucocorticoid receptor) antagonist, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual
- 35 syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment

disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
10 factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of
15 a GR antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GR antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GR antagonist compound to augment the
25 therapeutic effect or to provide a faster onset of the therapeutic effect of said GR antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- According to a preferred embodiment, the invention relates to the use as described above,
30 wherein said GR antagonist compound is chosen from the group consisting of ORG 34517/34850 and mifepristone, preferably mifepristone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, mifepristone is to be administered in a daily dose of between 600 and 1200 mg of the active ingredient.
- 35 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a GR antagonist, preferably chosen from the group consisting of ORG

34517/34850 and mifepristone, preferably mifepristone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said GR antagonist is Mifepristone, preferably provided in a unitary dose of between 600 and 1200 mg of the active ingredient.

42: combination therapy with a histamine H3-receptor antagonist

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a histamine H₃-receptor antagonist, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized

in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a histamine H3-receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said histamine H3-receptor antagonist compound, further
5 characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said histamine H3-receptor antagonist compound is chosen from the
10 group of compounds consisting of ABT-834 and ABT-239, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a histamine H3-receptor antagonist, preferably chosen from the group consisting
15 of ABT-834 and ABT-239 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of Alzheimer
20 Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

25 43: combination therapy with a hormonal substance

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a hormonal substance, are chosen from the group of diseases or disorders consisting
30 of premenstrual syndrome and sexual and gender identity disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group
35 of diseases and disorders consisting of premenstrual syndrome and sexual and gender identity disorders, characterized in that pipamperon or said pharmaceutically acceptable

salt thereof is administered simultaneously with, separate from or prior to the administration of a hormonal substance to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said hormonal substance, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said hormonal substance is chosen from the group consisting of a testosterone transdermal spray, a testosterone gel, a female testosterone patch, synthetic conjugated estrogen A, methyltestosterone, a estrogens/methyltestosterone and a drosiperone/ethinyl estradiol composition, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said hormonal substance is synthetic conjugated estrogen A and is to be administered in a daily dose ranging between 0.075 and 0.6 mg of the active ingredient. More preferably, said hormonal substance is a drosiperone/ethinyl estradiol composition and is to be administered as a daily dose in tablets, preferably comprising 3mg drosiperone and 0.02 mg ethinyl estradiol of the active ingredients, respectively.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a hormonal substance, preferably chosen from the group consisting of a testosterone transdermal spray, a testosterone gel, a female testosterone patch, synthetic conjugated estrogen A, methyltestosterone, a estrogens/methyltestosterone and a drosiperone/ethinyl estradiol composition, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of premenstrual syndrome and sexual and gender identity disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said hormonal substance is synthetic conjugated estrogen A, preferably provided in a unitary dose of between 0.075 and 0.6 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said hormonal substance is a drosiperone/ethinyl estradiol composition, preferably provided in tablets comprising a unitary dose of 3mg drosiperone and 0.02 mg ethinyl estradiol of the active ingredients, respectively.

5 44: combination therapy with a compound which increases brain concentrations of 5-HT

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which increases brain concentrations of 5-HT (serotonin), are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases brain concentrations of 5-HT (serotonin) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases brain concentrations of 5-HT (serotonin), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases brain concentrations of 5-HT (serotonin) to

35

augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases brain concentrations of 5-HT (serotonin), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which increases brain concentrations of 5-HT (serotonin) is chosen from the group consisting of triptosine, tramadol, SP 186, PMD 145 and KW 6055, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

10

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which increases brain concentrations of 5-HT (serotonin), preferably chosen from the group consisting of triptosine, tramadol, SP 186, PMD 145 and KW 6055, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15

20

25

45: combination therapy with a compound which increases insulin sensitivity

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which increases insulin sensitivity, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

30

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases insulin sensitivity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases Insulin sensitivity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said compound which increases insulin sensitivity is rosiglitazone maleate, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

20

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which increases insulin sensitivity, preferably rosiglitazone maleate or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

30

46: combination therapy with a compound inhibiting the mixed lineage kinase family

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy

35

with a compound which is an inhibitor of the mixed lineage kinase family is Parkinson Disease.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is an inhibitor of the mixed lineage kinase family to augment the therapeutic effect or to provide a faster onset of the
10 therapeutic effect of said compound which is an inhibitor of the mixed lineage kinase family, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 According to a preferred embodiment, the invention relates to the use as described above, wherein said compound which is an inhibitor of the mixed lineage kinase family is CEP-1347 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

20 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is an inhibitor of the mixed lineage kinase family, preferably CEP-1347 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

25 47: combination therapy with an interleukin-1 beta converting enzyme inhibitor compound
The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an interleukin-1 beta converting enzyme inhibitor compound, is a pain disorder.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an interleukin-1 beta converting enzyme inhibitor
35 compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said interleukin-1 beta converting enzyme inhibitor compound, further

characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above,
5 wherein said interleukin-1 beta converting enzyme inhibitor is pralnacasan or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an interleukin-1 beta converting enzyme inhibitor, preferably pralnacasan or a pro-
10 drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a pain disorder,.

48: combination therapy with a levodopa/decarboxylase inhibitor compound

15 The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a levodopa/decarboxylase inhibitor compound, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically
20 acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a levodopa/decarboxylase inhibitor compound to
25 augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said levodopa/decarboxylase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above,
30 wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa, levodopa/benserazide, etilevodopa/carbidopa or etilevodopa/benserazide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. According to a further preferred embodiment, the invention relates to the use as described above, wherein said
35 levodopa/decarboxylase inhibitor compound is (eti)levodopa/carbidopa, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof in combination with entacapone, which is an inhibitor of catechol-O-methyltransferase (COMT), or a pro-

drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa and is to be administered in a dose ranging between 2000 mg/ 50 mg and 100 mg/ 10 mg of the active ingredients. Preferably said entacapone is to be administered in a dose ranging
5 between 1000 mg/ 50 mg, more preferably between 500 mg/ 100 mg, and most preferably 200 mg of the active ingredients per day.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a levodopa/decarboxylase inhibitor compound, preferably levodopa/carbidopa,
10 levodopa/ benserazide, etilevodopa/carbidopa or etilevodopa/benserazide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease. The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a levodopa/decarboxylase
15 inhibitor compound, preferably is (eti)levodopa/carbidopa, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof in combination with entacapone, which is an inhibitor of catechol-O-methyltransferase (COMT), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the
20 underlying emotion dysregulation of Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa, preferably provided in a unitary dose of between 100 mg and 10 mg of the active
25 ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa or
30 etilevodopa/ carbidopa in combination with entacapone, of which the latter is preferably provided in a unitary dose of between 500 mg and 100 mg of the active ingredient.

49: combination therapy with a lipid-DNA complex

The mental disorder which can be treated using compounds having a high selective
35 affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a lipid-DNA complex, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said
5 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of lipid-DNA complex to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said lipid-DNA complex, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

According to a preferred embodiment, the invention relates to the use as described above, wherein said lipid-DNA complex is GR213487B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

15 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a lipid-DNA complex, preferably GR213487B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

20

50: combination therapy with a monoamine oxidase (MAO) reuptake inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase (MAO) reuptake inhibitor, are chosen from the group of
25 diseases or disorders consisting of substance related disorders and attention-deficit disorders (ADHD).

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
30 emotion dysregulation of non-cognitive mental disease or disorder which are substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase (MAO) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
35 monoamine oxidase (MAO) reuptake inhibitor compound, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of attention-deficit disorders (ADHD), characterized in that
10 pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase (MAO) reuptake inhibitor compound to augment the therapeutic effect or to provide a
15 faster onset of the therapeutic effect of said monoamine oxidase (MAO) reuptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 According to a preferred embodiment, the invention relates to the uses as described above, wherein said monoamine oxidase (MAO) reuptake inhibitor compound is NS 2359 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

20 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase (MAO) reuptake inhibitor, preferably NS 2359 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of substance related disorders and attention-deficit disorders
25 (ADHD).

51: combination therapy with a MAO-A and a MAO-B reuptake inhibitor

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy
30 with a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor, wherein said disorders are attention-deficit disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
35 emotion dysregulation of attention-deficit disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with,

separate from or prior to the administration of a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound, further
5 characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-
10 B) reuptake inhibitor compound is SPD473 or a pro-drug or an active metabolite thereof, or a pharma-ceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake
15 inhibitor, preferably SPD473 or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of attention-deficit disorders.

52: combination therapy with a monoamine oxidase B (MAO-B) inhibitor

20 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase B (MAO-B) inhibitor, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual
syndrome, somatoform disorders (excluding pain disorders), factitious disorders,
25 dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorder and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically
30 acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
factitious disorders, dissociative disorders, adjustment disorders, impulse control
35 disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, characterized in that pipamperon or said

pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that
5 pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
10 emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of
15 said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
20 emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of
25 said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said monoamine oxidase B (MAO-B) inhibitor compound is chosen from
30 the group consisting of selegiline, rasagiline (TVP-1012) and EmSam (transdermal selegiline), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said monoamine oxidase B (MAO-B) inhibitor is selegiline and is to be administered in a daily dose ranging between 5 and 10 mg of the active ingredient. More preferably, said monoamine oxidase B (MAO-B) inhibitor is rasagiline
35 (TVP-1012) and is to be administered in a daily dose ranging between 1 and 2 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase B (MAO-B) inhibitor, preferably chosen from the group consisting of selegiline, rasagiline (TVP-1012) and EmSam (transdermal selegiline), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorder and Parkinson Disease.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said monoamine oxidase B (MAO-B) inhibitor is selegiline, preferably provided in a unitary dose of between 5 and 10 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said monoamine oxidase B (MAO-B) inhibitor is rasagiline (TVP-1012), preferably provided in a unitary dose of between 1 and 2 mg of the active ingredient.

53: combination therapy with a monoamine oxidase B (MAO-B) reuptake inhibitor

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a monoamine oxidase B (MAO-B) reuptake inhibitor, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) reuptake inhibitor to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) reuptake inhibitor, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 According to a preferred embodiment, the invention relates to the use as described above, wherein said monoamine oxidase B (MAO-B) reuptake inhibitor is safinamide or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a monoamine oxidase B (MAO-B) reuptake inhibitor, preferably safinamide or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

54: combination therapy with a melanocortin-4 (MC4) receptor antagonist compound

15 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a melanocortin-4 (MC4) receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious
20 disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

25 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
30 factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of
35 a melanocortin-4 (MC4) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanocortin-4 (MC4) receptor

antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melanocortin-4 (MC4) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanocortin-4 (MC4) receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said melanocortin-4 (MC4) receptor antagonist compound is MCL0129, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a melanocortin-4 (MC4) receptor antagonist compound, preferably MCL0129 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

55: combination therapy with a MCH receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a melanin concentrating hormone (MCH) receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity

disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
10 factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of
15 a melanin concentrating hormone (MCH) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanin concentrating hormone (MCH) receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
25 from or prior to the administration of a melanin concentrating hormone (MCH) receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melanin concentrating hormone (MCH) receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 According to a preferred embodiment, the invention relates to the uses as described above, wherein said melanin concentrating hormone (MCH) receptor antagonist compound is SNAP-7941 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

35

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a melanin concentrating hormone (MCH) receptor antagonist compound, preferably SNAP-7941 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

56: combination therapy with a melatonin receptor (MT) agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a melatonin receptor (MT) agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melatonin receptor (MT) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melatonin receptor (MT) agonist

compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melatonin receptor (MT) agonist compound to
10 augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melatonin receptor (MT) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 According to a preferred embodiment, the invention relates to the uses as described above, wherein said melatonin receptor (MT) agonist compound is chosen from the group consisting of ramelteon and agomelatine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said melatonin receptor (MT) agonist compound is agomelatine and is to be administered in a daily dose ranging between 25 and 50 mg of the active ingredient.

20 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a melatonin receptor (MT) agonist compound, preferably ramelteon or agomelatine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or
25 sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems
30 related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient
35 and wherein said melatonin receptor (MT) agonist compound is agomelatine, preferably provided in a unitary dose of between 25 and 50 mg of the active ingredient.

57: combination therapy with a metabotropic glutamate receptor (MgluR) agonist compound

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a metabotropic glutamate receptor (MgluR) agonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, 10 bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying 15 emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, 20 characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a metabotropic glutamate receptor (MgluR) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said metabotropic glutamate receptor (MgluR) agonist compound, further characterized in that pipamperon is to be 25 administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the ~~preparation of~~ a medicament for treating the underlying 30 emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a metabotropic glutamate receptor (MgluR) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said metabotropic glutamate receptor (MgluR) agonist compound, further 35 characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said metabotropic glutamate receptor (MgluR) agonist compound is PRE703 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a metabotropic glutamate receptor (MgluR) agonist, preferably PRE703 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

58: combination therapy with a compound mimicking the effect of nerve growth factor (NGF)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which mimics the effect of nerve growth factor (NGF), are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic

disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which mimics the effect of nerve growth factor (NGF) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which mimics the effect of nerve growth factor (NGF), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which mimics the effect of nerve growth factor (NGF) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which mimics the effect of nerve growth factor (NGF), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which mimics the effect of nerve growth factor (NGF) is xaliproden or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said compound which mimics the effect of nerve growth factor (NGF) is xaliproden and is to be administered in a daily dose ranging between 1 and 2 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which mimics the effect of nerve growth factor (NGF), preferably xaliproden or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic

disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

- 5 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said compound which mimics the effect of nerve growth factor (NGF) is xaliproden, preferably provided in a unitary dose of between 1 and 2 mg of the active ingredient.

10

59: combination therapy with a muscarinic receptor partial agonist compound

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a muscarinic receptor partial agonist compound, are chosen from the group of
- 15 diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic
- 20 disorder, mild cognitive impairment disorder and other cognitive disorders.

- The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of
- 25 diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic
- 30 disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a muscarinic receptor partial agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said muscarinic receptor partial agonist compound, further
- 35 characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said muscarinic receptor partial agonist compound is sevimeline or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

5

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a muscarinic receptor partial agonist compound, preferably sevimeline or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the

10 underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general

15 medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

60: combination therapy with a selective nor-adrenaline re-uptake inhibitor (NARI) compound

20 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective nor-adrenaline re-uptake inhibitor (NARI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial

25 behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

30 emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is

35 administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline receptor inhibitor (NARI) compound to augment the therapeutic

effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline re-uptake inhibitor (NARI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
10 from or prior to the administration of a selective nor-adrenaline re-uptake inhibitor (NARI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline re-uptake inhibitor (NARI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective nor-adrenaline re-uptake inhibitor (NARI) compound is chosen from the group consisting of reboxetine, atomoxetine hydrochloride, A 75200, 155U88, (S)-A 75200, tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, viloxazine, tomoxetine, duloxetine, venlafaxine and milnacipran or a pro-drug or an active metabolite thereof, or a
20 pharmaceutically acceptable salt thereof. More preferably, said selective nor-adrenaline re-uptake inhibitor (NARI) compound is reboxetine and is to be administered in a daily dose ranging between 8 and 12 mg of the active ingredient. More preferably, said selective nor-adrenaline re-uptake inhibitor (NARI) compound is atomoxetine hydrochloride and is to be administered in a daily dose ranging between 40 and 100 mg of the active ingredient.

25

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective nor-adrenaline re-uptake inhibitor (NARI) compound, preferably
30 chosen from the group consisting of reboxetine, atomoxetine hydrochloride, A 75200, 155U88, (S)-A 75200, tandamine, pirandamine, ciclazindol, fluparoxan, lortalamine, talsupram, talopram, prindamine, nomifensine, viloxazine, tomoxetine, duloxetine, venlafaxine and milnacipran, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous,
35 separate or sequential use for treating the underlying emotion dysregulation of mental

disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

5

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective nor-adrenaline re-uptake inhibitor (NARI) compound is reboxetine, preferably provided in a unitary dose of between 8 and 12 mg of the active

10

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective nor-adrenaline re-uptake inhibitor (NARI) compound is atomoxetine hydrochloride, preferably provided in a unitary dose of between 40 and 100 mg of the active ingredient.

15

61: combination therapy with a NaSSA compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a noradrenergic/specific serotonergic antidepressant (NaSSA) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

20

25

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse

30

35

or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a noradrenergic/specific serotonergic antidepressant (NaSSA) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said
5 noradrenergic/specific serotonergic antidepressant (NaSSA) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically
10 acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a noradrenergic/specific serotonergic antidepressant (NaSSA) compound to augment the therapeutic effect or to provide a faster onset of the
15 therapeutic effect of said noradrenergic/specific serotonergic antidepressant (NaSSA) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described
20 above, wherein said noradrenergic/specific serotonergic antidepressant (NaSSA) compound is ORG 4420 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon,
25 and (b) a noradrenergic/specific serotonergic antidepressant (NaSSA) compound, preferably ORG 4420 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders,
30 eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

62: combination therapy with a selective NDRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, are

5 chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head

10 trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial

20 behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective

25 nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said

30 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset

35 of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor

(NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound is GW353162 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound is GW353162 and is to be administered in a daily dose ranging between 20 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, preferably GW353162 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia,

vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound is GW353162, preferably provided in a unitary dose of between 20 and 60 mg of the active ingredient.

63: combination therapy with a compound which is a neuroimmunophilin ligand

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which is a neuroimmunophilin ligand, is Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a neuroimmunophilin ligand to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a neuroimmunophilin ligand, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said a compound which is a neuroimmunophilin ligand is GPI.1485 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said a compound which is a neuroimmunophilin ligand is GPI 1485 and is to be administered in a daily dose ranging between 200 and 1000 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is a neuroimmunophilin ligand, preferably GPI 1485 or a pro-

drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

- 5 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said a compound which is a neuroimmunophilin ligand is GPI 1485, preferably provided in a unitary dose of between 200 and 1000 mg of the active ingredient.

10

64: combination therapy with a neuromodulator compound

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neuromodulator compound, is Parkinson Disease.

15

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neuromodulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neuromodulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20

- 25 According to a preferred embodiment, the invention relates to the use as described above, wherein said neuromodulator compound is adenosine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

30

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neuromodulator compound, preferably adenosine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

65: combination therapy with a neurotensin receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neurotensin receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor antagonist compound, further characterized in that pipamperon is to

be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said neurotensin receptor antagonist compound is SR 48692 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said neurotensin receptor antagonist compound is SR 48692 and is to be administered in a daily dose ranging between 90 and 300 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neurotensin receptor antagonist compound, preferably SR 48692 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said neurotensin receptor antagonist compound is SR 48692, preferably provided in a unitary dose of between 90 and 300 mg of the active ingredient.

66: combination therapy with nerve growth factor (NGF) gene therapy

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with nerve growth factor (NGF) gene therapy, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from nerve growth factor (NGF) gene therapy, to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nerve growth factor (NGF) gene therapy, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to nerve growth factor (NGF) gene therapy, to augment the therapeutic effect or to provide a faster onset of nerve growth factor (NGF) gene therapy, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound useful in nerve growth factor (NGF) gene therapy, preferably xaliproden or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

It should be understood that "nerve growth factor gene therapy" is well known in the art, and the compounds, for instance nucleic acids used in nerve growth factor gene therapy are well described (see e.g. Tuszynski *et al.*, (2002) Journal of Molecular Neuroscience Volume 19, Issue 1-2, pps. 207-208).

67: combination therapy with a nicotinic acetylcholine receptor antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a nicotinic acetylcholine receptor antagonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is

115

administered simultaneously with, separate from or prior to the administration of a nicotinic acetylcholine receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic acetylcholine receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a nicotinic acetylcholine receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic acetylcholine receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said nicotinic acetylcholine receptor antagonist compound is SEP174559 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a nicotinic acetylcholine receptor antagonist compound, preferably SEP174559 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of anxiety disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

68: combination therapy with a nicotinic receptor agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a nicotinic receptor agonist compound, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia,

vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a nicotinic receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said nicotinic receptor agonist compound is ABT-089, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said nicotinic receptor agonist compound is ABT-089 and is to be administered in a daily dose ranging between 4 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a nicotinic receptor agonist compound, preferably ABT-089 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease,

dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

- 5 The invention also relates to a pharmaceutical composition as described above, wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said nicotinic receptor agonist compound is ABT-089, preferably provided in a unitary dose of between 4 and 40 mg of the active ingredient.

10 69: combination therapy with a neurokinin 2 receptor (NK2) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neurokinin 2 receptor (NK2) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders,
15 premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

20

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating
25 disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt
30 thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 2 receptor (NK2) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 2 receptor (NK2) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
5 from or prior to the administration of a neurokinin 2 receptor (NK2) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 2 receptor (NK2) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10 According to a preferred embodiment, the invention relates to the uses as described above, wherein said neurokinin 2 receptor (NK2) antagonist compound is saredutant or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said neurokinin 2 receptor (NK2) antagonist compound is saredutant and
15 is to be administered in a daily dose ranging between 25 and 200 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neurokinin 2 receptor (NK2) antagonist compound, preferably saredutant or a
20 pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders,
25 dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

30 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said neurokinin 2 receptor (NK2) antagonist compound is saredutant, preferably provided in a unitary dose of between 25 and 200 mg of the active ingredient.

70: combination therapy with a neurokinin 3 receptor (NK3) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a neurokinin 3 receptor (NK3) antagonist compound, are chosen from the group of

5 diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions,

10 malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

15 emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related

20 disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound

25 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound

35 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that

pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to
10 provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 According to a preferred embodiment, the invention relates to the uses as described above, wherein said neurokinin 3 receptor (NK3) antagonist compound is talnetant or osanetant, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said neurokinin 3 receptor (NK3) antagonist compound is talnetant and is to be administered in a daily dose ranging between 1.5 and 12 mg of the active ingredient.

20 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a neurokinin 3 receptor (NK3) antagonist compound, preferably talnetant or osanetant, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for
25 treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering,
30 antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

35 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

and wherein said neurokinin 3 receptor (NK3) antagonist compound is talnetant, preferably provided in a unitary dose of between 1.5 and 12 mg of the active ingredient.

71: combination therapy with an N-Methyl-D-aspartate (NMDA) antagonist compound

- 5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an N-Methyl-D-aspartate (NMDA) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious
- 10 disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to
- 15 Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.
- 20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
- 25 factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-
- 30 aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 35 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying

emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound; further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said N-Methyl-D-aspartate (NMDA) antagonist compound is chosen from the group consisting of SEP174559, memantine, delucemine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said N-Methyl-D-aspartate (NMDA) antagonist compound is memantine and is to be administered in a daily dose ranging between 5 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an N-Methyl-D-aspartate (NMDA) antagonist compound, preferably chosen from the group consisting of SEP174559, memantine, delucemine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying

emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said N-Methyl-D-aspartate (NMDA) antagonist compound is memantine, preferably provided in a unitary dose of between 5 and 40 mg of the active ingredient.

72: combination therapy with a non-steroidal anti-inflammatory drug

The mental disorder which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a non-steroidal anti-inflammatory drug, is a pain disorder or Alzheimer Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a non-steroidal anti-inflammatory drug to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said a non-steroidal anti-inflammatory drug, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disease, such as Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered

124

simultaneously with, separate from or prior to the administration of a non-steroidal anti-inflammatory drug to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said a non-steroidal anti-inflammatory drug, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said a non-steroidal anti-inflammatory drug is chosen from the group consisting of piroxicam, MX-1094, meloxicam and flurizan (pure R-enantiomer form of flurbiprofen), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a non-steroidal anti-inflammatory drug, preferably chosen from the group consisting of piroxicam, MX-1094, meloxicam and flurizan (pure R-enantiomer form of flurbiprofen), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a pain disorder or Alzheimer Disease.

73: combination therapy with an opoid antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an opoid antagonist compound, are substance related disorders.

It will be appreciated that the terms "opoid" and "opioid" may be used interchangeably.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a opoid antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said opoid antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said opoid antagonist compound is naltrexone, preferably as a depot formulation, more preferably in the form of microcapsules, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Preferably, said naltrexone is to be administered in the form of a depot, preferably a depot of microcapsules comprising a daily dose of between 192 and 384 mg.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a opoid antagonist, preferably naltrexone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of substance related disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said opoid antagonist compound is naltrexone, preferably provided in a unitary dose of between 192 and 384 mg of the active ingredient.

74: combination therapy with an opoid agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with an opoid agonist compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and

126

problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an opioid agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said opioid agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said opioid agonist compound is chosen from the group consisting of siramesine, E-5842 and cyclazocine, preferably siramesine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) an opioid agonist compound, preferably chosen from the group consisting of siramesine, E-5842 and cyclazocine, preferably siramesine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group consisting of anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

75: combination therapy with a phosphodiesterase-4 (PDE4) inhibitor compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a phosphodiesterase-4 (PDE4) inhibitor compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob

Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
10 factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of phosphodiesterase-4
15 (PDE4) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phosphodiesterase-4 (PDE4) inhibitor compound d
25 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 30 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head
35 trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic

- disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phosphodiesterase-4 (PDE4) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 10 According to a preferred embodiment, the invention relates to the uses as described above, wherein said phosphodiesterase-4 (PDE4) inhibitor compound is chosen from the group consisting of ND1251 and MEM 1917 (R1497), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.
- 15 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a phosphodiesterase-4 (PDE4) inhibitor antagonist compound, preferably chosen from the group consisting of ND1251 and MEM 1917 (R1497), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying
- 20 emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to
- 25 abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic
- 30 disorder, mild cognitive impairment disorder and other cognitive disorders.

76: combination therapy with a peptidic compound

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a peptidic compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome,
- 35

somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

10

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15

20

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a ~~peptidic~~ compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said ~~peptidic~~ compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group

35

of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said peptidic compound is chosen from the group consisting of secretin, PT-141, INN 00835 and beta sheet breaker peptide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said peptidic compound is secretin and is to be administered in a daily dose ranging between 0.2 and 0.4 mg/kg of the active ingredient. More preferably, said peptidic compound is INN 00835 and is to be administered in a daily dose ranging between 18 and 160 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a peptidic compound, preferably chosen from the group consisting of secretin, PT-141, INN 00835 and beta sheet breaker peptide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition,

substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

5 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said peptidic compound is secretin, preferably provided in a unitary dose of 0.2 and 0.4 mg/kg of the active ingredient.

10 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said peptidic compound is INN 00835, preferably provided in a unitary dose of 18 and 160 mg of the active ingredient.

77: combination therapy with a phospholipase A2 inhibitor compound

15 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a phospholipase A2 inhibitor compound which has caspase inhibitor activity, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

25

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the

30

35

administration of a phospholipase A2 inhibitor compound which has caspase inhibitor activity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phospholipase A2 inhibitor compound which has caspase inhibitor activity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phospholipase A2 inhibitor compound which has caspase inhibitor activity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phospholipase A2 inhibitor compound which has caspase inhibitor activity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phospholipase A2 inhibitor compound which has caspase inhibitor activity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phospholipase A2 inhibitor compound which has caspase inhibitor activity, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said phospholipase A2 inhibitor compound which has caspase inhibitor activity is chosen from the group consisting of LAX-101a, LAX-101b and LAX-101c, preferably LAX-101a, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a phospholipase A2 inhibitor compound which has caspase inhibitor activity, preferably chosen from the group consisting of LAX-101a, LAX-101b and LAX-101c, more

preferably LAX-101a, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

78: combination therapy with a compound which is a prodrug of uridine

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which is a prodrug of uridine, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a prodrug of uridine to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a prodrug of uridine, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
5 from or prior to the administration of a compound which is a prodrug of uridine to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a prodrug of uridine, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10 According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which is a prodrug of uridine is RG2133 (triacetyluridine) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

15 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is a prodrug of uridine, preferably RG2133 (triacetyluridine) or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the
20 underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational
25 problem, problems related to abuse or neglect and pain disorders.

79: combination therapy with prostaglandin E1 compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy
30 with prostaglandin E1 compound, are sexual and gender identity disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the
35 group of sexual and gender identity disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate

from or prior to the administration of a prostaglandin E1 compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said prostaglandin E1 compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the use as described above, wherein said prostaglandin E1 is alprostadil or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said prostaglandin E1 compound is alprostadil, preferably in the form of cream or gel, preferably a topical gel, and is to be administered in a daily dose ranging between 50 and 300 microgram per application of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a prostaglandin E1 compound, preferably alprostadil or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of sexual and gender identity disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said prostaglandin E1 compound is alprostadil, preferably provided in the form of a cream or gel, preferably a topical gel, wherein a unitary dose comprises between 50 and 300 microgram of the active ingredient per application.

80: combination therapy with a compound protecting dopaminergic and cholinergic neurons

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which protects dopaminergic and cholinergic neurons, are chosen from the group of diseases or disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic

disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which protects dopaminergic and cholinergic neurons to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which protects dopaminergic and cholinergic neurons, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which protects dopaminergic and cholinergic neurons to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which protects dopaminergic and cholinergic neurons, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which protects dopaminergic and cholinergic neurons is SR 57667 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which protects dopaminergic and cholinergic neurons, preferably SR 57667 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

81: combination therapy with a psychostimulant

- 15 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a psychostimulant, are chosen from the group of diseases or disorders consisting of sleep disorders, attention-deficit disorders and substance-related disorders.
- 20 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of sleep disorders, attention-deficit disorders and substance-related disorders, characterized in that pipamperon or said
- 25 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a psychostimulant to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said psychostimulant, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 30 According to a preferred embodiment, the invention relates to the uses as described above, wherein said psychostimulant is chosen from the group consisting of SPD 503, r-modafinil and modafinil, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said psychostimulant is SPE
- 35 503, more preferably said psychostimulant is modafinil and is to be administered in a daily dose ranging between 200 and 600 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a psychostimulant, preferably chosen from the group consisting of SPD 503, r-modafinil and modafinil, more preferably said SPC 503 or modafinil or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group consisting of sleep disorders, attention-deficit disorders and substance-related disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said psychostimulant is modafinil, preferably provided in a unitary dose of between 200 and 600 mg of the active ingredient.

82: combination therapy with a compound which is a Reversible Inhibitor of Mono-Amine oxydase A (RIMA)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is

administered simultaneously with, separate from or prior to the administration of a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), further
5 characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
10 emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which is a reversible inhibitor of mono-amine
15 oxydase A (RIMA), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said compound which is a reversible inhibitor of mono-amine oxydase A
20 (RIMA) is chosen from the group consisting of toloxatone, RS 8359, moclobemide, cimoxatone, caroxazone (F.I 6654) and befloxatone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) is befloxatone and is to be administered in a daily dose ranging between 2.5 and 20 mg of the active ingredient.

25 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), preferably chosen from the group consisting of toloxatone, RS 8359, moclobemide, cimoxatone, caroxazone (F.I 6654) and befloxatone, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for
30 simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual
35 and gender identity disorders, adjustment disorders, impulse control disorders, personality

disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

5 The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) is befloxadone, preferably provided in a unitary dose of between 2.5 and 20 mg of the active ingredient.

10 83: combination therapy with a compound which modulates SCT-11

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound which modulates SCT-11 (i.e. SCT-11 is a G protein-coupled receptor), are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

20

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which modulates SCT-11 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which modulates SCT-11, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

141

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
5 from or prior to the administration of a compound which modulates SCT-11 to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which modulates SCT-11, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

According to a preferred embodiment, the invention relates to the use as described above, wherein said compound which modulates SCT-11 is SNEC-2 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

15 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound which modulates SCT-11, preferably SNE-2 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group
20 consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

25

84: combination therapy with a serotonin/dopamine antagonist compound (SDA)

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a serotonin/dopamine antagonist compound (SDA), are chosen from the group of
30 diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem,
35 problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine antagonist compound,

further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described
5 above, wherein said serotonin/dopamine antagonist compound is chosen from the group consisting of zotepine, ziprasidone, SM-13496, SL 91.0177, sertindole, S-18327, risperidone, quetiapine fumarate (preferably sustained release formulation), quetiapine fumarate (preferably granules), quetiapine, perospirone, paliperidone, olanzapine, ocaperidone, LU 31-131, iloperidone, clozapine, BSF-190555, blonanserin, bifeprunox,
10 asenapine and aripiprazole, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Even more preferably, said serotonin/dopamine antagonist compound is chosen from the group consisting of SL 91.0177, sertindole, perospirone, paliperidone, blonanserin, bifeprunox and asenapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably,
15 said serotonin/dopamine antagonist compound is sertindole and is to be administered in a daily dose ranging between 12 and 24 mg of the active ingredient. More preferably, said serotonin/dopamine antagonist compound is paliperidone and is to be administered in a daily dose ranging between 3 and 15 mg of the active ingredient. More preferably, said serotonin/dopamine antagonist compound is asenapine and is to be administered in a
20 daily dose ranging between 2.5 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a serotonin/dopamine antagonist compound, preferably chosen from the group consisting of SL 91.0177, sertindole, perospirone, paliperidone, blonanserin, bifeprunox
25 and asenapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders,
30 dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

35

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said serotonin/dopamine antagonist compound is sertindole, preferably provided in a unitary dose of between 12 and 24 mg of the active ingredient.

5

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said serotonin/dopamine antagonist compound is paliperidone, preferably provided in a unitary dose of between 3 and 15 mg of the active ingredient.

10

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said serotonin/dopamine antagonist compound is asenapine, preferably provided in a unitary dose of between 2.5 and 20 mg of the active ingredient.

15

85: combination therapy with a selective SDRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin and dopamine re-uptake inhibitor (SDRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder and other cognitive disorders.

30

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),

35

factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5 According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin and dopamine reuptake inhibitor (SDRI) compound is bazineprine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective serotonin and dopamine reuptake inhibitor (SDRI) compound, preferably bazineprine or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, 15 eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting 20 dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

25

86: combination therapy with a second messenger beta agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy 30 with a second messenger beta agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and 35 pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a second messenger beta agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said second messenger beta agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a second messenger beta agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said second messenger beta agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said second messenger beta agonist compound is chosen from the group consisting of SR 57227, rolipram and eplivanserin, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said second messenger beta agonist compound is ~~rolipram~~ and is to be administered in a daily dose ranging between 1.5 and 3 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a second messenger beta agonist compound, preferably chosen from the group consisting of SR 57227, rolipram and eplivanserin or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for

simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said second messenger beta agonist compound is rolipram, preferably provided in a unitary dose of between 1.5 and 3 mg of the active ingredient.

87: combination therapy with a secretin pancreatic hormone

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a secretin pancreatic hormone, are chosen from the group of diseases or disorders consisting of anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the

administration of a secretin pancreatic hormone to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said secretin pancreatic hormone, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a secretin pancreatic hormone to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said secretin pancreatic hormone, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a secretin pancreatic hormone to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said secretin pancreatic hormone, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20

25 According to a preferred embodiment, the invention relates to the uses as described above, wherein said secretin pancreatic hormone is RG1068 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a secretin pancreatic hormone, preferably RG1068, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive

30

35

behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

5

88: combination therapy with a sigma receptor agonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a sigma receptor agonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor agonist compound to augment the

35

therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

- 5 According to a preferred embodiment, the invention relates to the uses as described above, wherein said sigma receptor agonist compound is VPI-013 (also known as OPC-14523) or PRX-00023, preferably VPI-013 (also known as OPC-14523), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.
- 10 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a sigma receptor agonist compound, preferably VPI-013 (also known as OPC-14523) or PRX-00023, preferably VPI-013 (also known as OPC-14523), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying
- 15 emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, personality disorders, bereavement,
- 20 occupational problem, problems related to abuse or neglect and pain disorders.

89: combination therapy with a sigma receptor antagonist compound

- The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy
- 25 with a sigma receptor antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development
- 30 disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

- The present invention thus relates to the use of pipamperon or a pharmaceutically
- 35 acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the

group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect,, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sigma receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sigma receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said sigma receptor antagonist compound is chosen from the group consisting of SR 31742 and EMD 68843, or a pro-drug or an active metabolite thereof, or

a pharmaceutically acceptable salt thereof. More preferably, said sigma receptor antagonist compound is EMD 68843 and is to be administered in a daily dose ranging between 5 and 40 mg of the active ingredient.

- 5 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a sigma receptor antagonist compound, preferably chosen from the group consisting of SR 31742 and EMD 68843, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental
- 10 disease or disorder which is chosen from the group consisting mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-
- 15 related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect, pain disorders and delirium.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient

20 and wherein said sigma receptor antagonist compound is EMD 68843, preferably provided in a unitary dose of between 5 and 40 mg of the active ingredient.

90: combination therapy with a selective SNDRI compound

- 25 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep
- 30 disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease,
- 35 dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition,

substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 1.5 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders consisting of delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to

head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized
5 in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further
10 characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor
15 (SNDRI) compound is selected from the group consisting of NS 2330; McN 5652; DOV 216,303 and DOV 21,947; more preferably NS 2330 or DOV 216,303; or a pro-drug or an active metabolite thereof; or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon,
20 and (b) a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, preferably selected from the group consisting of NS 2330; McN 5652; DOV 216,303 and DOV 21,947, more preferably NS 2330 or DOV 216,303, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying
25 emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders,
30 delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and
35 other cognitive disorders.

91: combination therapy with a selective SNRI compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and nor-adrenaline re-uptake inhibitor

(SNRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described
5 above, wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is selected from the group consisting of venlafaxine, tomoxetine, tandamine, talsupram, talopram, nefazodone, milnacipran, LY 113.821, duloxetine, desvenlafaxine and amoxapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Even more preferably, said selective serotonin and nor-adrenaline
10 re-uptake inhibitor (SNRI) compound is chosen from the group consisting of venlafaxine, tomoxetine, milnacipran, duloxetine and desvenlafaxine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is venlafaxine and is to be administered in a daily dose ranging between 75 and 300 mg of the active
15 ingredient. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is tomoxetine and is to be administered in a daily dose ranging between 0.475 and 3.8 mg/kg of the active ingredient. More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is milnacipran and is to be administered in a daily dose ranging between 50 and 200 mg of the active ingredient.
20 More preferably, said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is duloxetine and is to be administered in a daily dose ranging between 40 and 60 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon,
25 and (b) a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, preferably selected from the group consisting of venlafaxine, tomoxetine, tandamine, talsupram, talopram, nefazodone, milnacipran, LY 113.821, duloxetine, desvenlafaxine and amoxapine, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or
30 sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related
35 disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is venlafaxine, preferably provided in a unitary dose of between 75 and 300 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is tomoxetine, preferably provided in a unitary dose of between 0.475 and 3.8 mg/kg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is milnacipran, preferably provided in a unitary dose of between 50 and 200 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound is duloxetine, preferably provided in a unitary dose of between 40 and 60 mg of the active ingredient.

92: combination therapy with a selective serotonin re-uptake inhibitor (SSRI) compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a selective serotonin re-uptake inhibitor (SSRI) compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin re-uptake inhibitor (SSRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin re-uptake inhibitor (SSRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin re-uptake inhibitor (SSRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin re-uptake inhibitor (SSRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said selective serotonin re-uptake inhibitor (SSRI) compound is selected from the group consisting of YM 992, VPI-013 (also known as OPC-14523), sertraline, paroxetine, LY 214.281, LU AA-21-004, -Lu-35-138, litoxetine, ifoxetine, fluvoxamine (controlled release formulation), fluvoxamine, fluoxetine, femoxetine, escitalopram, EMD 68843, cyanodothepine, citalopram, venlafaxine, milnacipran, duloxetine, cericlamine and ademethionine (preferably s-adenosylmethionine), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. Even more preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is chosen from the group consisting of litoxetine, fluvoxamine (controlled release formulation) and escitalopram, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

More preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is fluvoxamine (controlled release formulation) and is to be administered in a daily dose ranging between 100 and 300 mg of the active ingredient. More preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is escitalopram and is to be administered in a daily dose ranging between 10 and 20 mg of the active ingredient. More preferably, said selective serotonin re-uptake inhibitor (SSRI) compound is citalopram and is to be administered in a daily dose ranging between 10 and 40 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a selective serotonin re-uptake inhibitor (SSRI) compound, preferably selected from the group consisting of YM 992, VPI-013 (also known as OPC-14523), sertraline, paroxetine, LY 214.281, LU AA 21-004, Lu 35-138, litoxetine, ifoxetine, fluvoxamine (controlled release formulation), fluvoxamine, fluoxetine, femoxetine, escitalopram, EMD 68843, cyanodothepine, citalopram, venlafaxine, milnacipran, duloxetine, cericlamine and ademethionine (preferably s-adenosylmethionine), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin re-uptake inhibitor (SSRI) compound is fluvoxamine (controlled release formulation), preferably provided in a unitary dose of between 100 and 300 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin re-uptake inhibitor (SSRI) compound is escitalopram, preferably provided in a unitary dose of between 10 and 20 mg of the active ingredient.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said selective serotonin re-uptake inhibitor (SSRI) compound is citalopram, preferably provided in a unitary dose of between 10 and 40 mg of the active ingredient.

5

Citalopram or citalopram hydrobromide is a selective serotonin (5-hydroxytryptamine / 5-HT) re-uptake inhibitor (SSRI) and is the conventional name given for the compound of the formula (RS)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalanecarbonitrile-hydro-bromide. According to an embodiment, a daily doses of active ingredient of SSRI, preferably citalopram, ranges between 10 and 40 mg per day. Preferably, daily doses of active ingredient ranging between 20 and 30 mg per day are administered. More preferably, a daily dose of 10, 15, 20, 25, 30, 35 or 40 mg per day is administered.

10

Fluvoxamine or fluvoxamine maleate (luvox, fevarin) is a selective serotonin (5-HT) re-uptake inhibitor (SSRI) belonging to a new chemical series, the 2-aminoethyl oxime ethers of aralkylketones. It is chemically unrelated to other SSRIs and clomipramine. It is chemically designated as 5-methoxy-4'-(trifluoromethyl) valerophenone (E)-O-(2-aminoethyl) oxime maleate (1:1).

15

According to an embodiment, a daily dose of active ingredient of fluvoxamine in a controlled release mode ranges between 100 and 300 mg per day. Preferably, daily doses of active ingredient ranging between 150 and 200 mg per day are administered in a controlled release mode. More preferably, a daily dose of 100, 150, 200, 250 or 300 mg per day is administered by controlled release.

20

93: combination therapy with a substance P receptor (NK1) antagonist compound

25

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a substance P receptor (NK1) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

30

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a substance P receptor (NK1) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said substance P receptor (NK1) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a substance P receptor (NK1) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said substance P receptor (NK1) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said substance P receptor (NK1) antagonist compound is chosen from the group consisting of vestipitant, TAK-637, R673, GW823296, GW679769, GW597599, CP-122.721, aprepitant, 823296 and 679769, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said substance P receptor (NK1) antagonist compound is aprepitant and is to be administered in a daily dose ranging between 40 and 160 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a substance P receptor (NK1) antagonist compound, preferably chosen from the group consisting of vestipitant, TAK-637, R673, GW823296, GW679769, GW597599, CP-

163

122.721, aprepitant, 823296 and 679769, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said substance P receptor (NK1) antagonist compound is aprepitant, preferably provided in a unitary dose of between 40 and 160 mg of the active ingredient.

94: combination therapy with a sulfonamide compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a sulfonamide compound, are chosen from the group of diseases or disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering,

antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sulfonamide compound to augment the therapeutic effect or to provide
5 a faster onset of the therapeutic effect of said sulfonamide compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically
10 acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sulfonamide compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sulfonamide compound,
15 further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
20 emotion dysregulation of a cognitive mental disease or disorder which is delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a sulfonamide compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said sulfonamide compound, further characterized in that
25 pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses as described above, wherein said sulfonamide compound is zonisamide or a pro-drug or an active
30 metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said sulfonamide compound is zonisamide and is to be administered in a daily dose ranging between 100 and 600 mg of the active ingredient.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon,
35 and (b) a sulfonamide compound, preferably zonisamide, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined

preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium.

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said sulfonamide compound is zonisamide, preferably provided in a unitary dose of between 100 and 600 mg of the active ingredient.

95: combination therapy with a tachykinin antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a tachykinin antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of

166

a tachykinin antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said tachykinin antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
10 from or prior to the administration of a tachykinin antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said tachykinin antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15 According to a preferred embodiment, the invention relates to the use as described above, wherein said tachykinin antagonist compound is SR 48968 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon,
20 and (b) a tachykinin antagonist compound, preferably SR 48968 or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome,
25 somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

30

96: combination therapy with a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493

35

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, are chosen from the group of

diseases or disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, phase of life problem, academic problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnestic disorders due to a general medical condition, substance-induced persisting amnestic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, phase of life problem, academic problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

168

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a compound selected from the group consisting of R228060 (YKP-10A), palanpanel, ORG 39479/PH80, ORG 34167, DP 543 and CJ-017.493, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined
5 preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders,
10 adjustment disorders, impulse control disorders, pervasive development disorders, attention-deficit disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, phase of life problem, academic problem, problems related to abuse or neglect, pain disorders,
15 delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other
20 cognitive disorders and Parkinson Disease.

97: combination therapy with a vasopressin 1B receptor (V1B) antagonist compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy
25 with a vasopressin 1B receptor (V1B) antagonist compound, are chosen from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality
30 disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying
35 emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating

disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders),
factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep
disorders, adjustment disorders, impulse control disorders, substance-related disorders,
personality disorders, bereavement, occupational problem and problems related to abuse
5 or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt
thereof is administered simultaneously with, separate from or prior to the administration of
a vasopressin 1B receptor (V1B) antagonist compound to augment the therapeutic effect
or to provide a faster onset of the therapeutic effect of said vasopressin 1B receptor (V1B)
antagonist compound, further characterized in that pipamperon is to be administered to a
10 patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically
acceptable salt thereof for the preparation of a medicament for treating the underlying
emotion dysregulation of pain disorders, characterized in that pipamperon or said
15 pharmaceutically acceptable salt thereof is administered simultaneously with, separate
from or prior to the administration of a vasopressin 1B receptor (V1B) antagonist
compound to augment the therapeutic effect or to provide a faster onset of the therapeutic
effect of said vasopressin 1B receptor (V1B) antagonist compound, further characterized
in that pipamperon is to be administered to a patient in a daily dose ranging between 5
20 and 15 mg of the active ingredient.

According to a preferred embodiment, the invention relates to the uses, as described
above, wherein said vasopressin 1B receptor (V1B) antagonist compound is SSR149415
or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt
25 thereof.

The invention also relates to a pharmaceutical composition comprising (a) pipamperon,
and (b) a vasopressin 1B receptor (V1B) antagonist compound, preferably SSR149415 or
a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof,
30 as a combined preparation for simultaneous, separate or sequential use for treating the
underlying emotion dysregulation of a mental disease or disorder which is chosen from
the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual
syndrome, somatoform disorders (excluding pain disorders), factitious disorders,
dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment
35 disorders, impulse control disorders, substance-related disorders, personality disorders,

bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

98: combination therapy with a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound

5 The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a voltage-gated calcium channel alpha(2)delta subunit modulator compound, are
10 disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating
20 disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt
25 thereof is administered simultaneously with, separate from or prior to the administration of a voltage-gated calcium channel alpha(2)delta subunit modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said voltage-gated calcium channel alpha(2)delta subunit modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5
30 and 15 mg of the active ingredient.

The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of pain disorders, characterized in that pipamperon or said
35 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a voltage-gated calcium channel alpha(2)delta

subunit modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

According to a preferred embodiment, the invention relates to the uses as described above, wherein said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound is pregabalin or PD-200,390; or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof. More preferably, said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound is pregabalin, and is to be administered in a daily dose ranging between 50 and 600 mg of the active ingredient.

10

The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) a voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound, preferably pregabalin or PD-200,390; or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a mental disease or disorder which is chosen from the group consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders.

15

20

The invention also relates to a pharmaceutical composition as described above wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said voltage-gated calcium channel $\alpha(2)\delta$ subunit modulator compound is pregabalin, preferably provided in a unitary dose of between 50 and 600 mg of the active ingredient.

25

30

99: combination therapy with a vomeropherin compound

The mental disorders which can be treated using compounds having a high selective affinity for the 5-HT_{2A} and D₄ receptor, for instance pipamperon, in a combination therapy with a vomeropherin compound, are chosen from the group of diseases or disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual

35

and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

- 5 The present invention thus relates to the use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of vomeropherin compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said vomeropherin compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 20 According to a preferred embodiment, the invention relates to the use as described above, wherein said vomeropherin compound is PH94B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

- 25 The invention also relates to a pharmaceutical composition comprising (a) pipamperon, and (b) vomeropherin compound, preferably PH94B or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder which is chosen from the group consisting of anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect.

- 35 Also, the invention relates in particular to the use as described before, wherein said second compound is chosen from the group consisting of fluvoxamine controlled release,

phenserine tartrate, atomoxetine hydrochloride, bupropion (controlled-release formulation), ropinirole HCL (controlled-release formulation), INN 00835, galantamine (extended release formulation), paliperidone, tomoxetine, aprepitant, rivastigmine tartrate, ORG 34517/34850, sunepitron, sumanirole, milnacipran, idazoxan, xaliprodin, SR 58611, 5 befloxadone, litoxetine, tianeptine, agomelatine, SPD 503, flesinoxan, bifeprunox, ramelteon, etilevodopa, rasagiline (TVP-1012) and desvenlafaxine.

Also, the invention relates in particular to the use as described before, wherein said second compound is chosen from the group consisting of galantamine (extended release formulation), R121919, risperidone, paliperidone and R228060 (YKP-10A).

10 The disclosure of all patents, publications (including published patent publications), and database accession numbers and depository accession numbers referenced in this specification are specifically incorporated herein by reference in their entirety to the same extent as if each such individual patent, publication, and database accession number, and 15 depository accession number were specifically and individually indicated to be incorporated by reference.

The invention, now being generally described, will be more readily understood by reference to the following tables and examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention and are not intended to limit the invention.

Short description of the Tables and Figures

Table 1: In Table 1, the pKi values of test compounds are given for each of the dopamine receptors, 5HT receptors, adrenergic receptors and the histamine1 receptor.

5

Table 2: Set-up of a clinical trial comprising for treatment groups.

Table 3: Overview of a placebo, active and period controlled clinical trial in a fore-going pipamperon - citalopram treatment in Major Depressive Disorder.

10

Table 4: POC process for major depressive disorder.

Table 5: Summary of diseases and disorders relative to known psycho-tropics.

15

Table 6: Overview of Pharmacological grouping, indicating pharmacological profile numbering (column 2), pharmacological profile (column 3), main indication(s) (column 4), name of the compound (column 4), the dose range (column 5), and the company producing or selling said compound (column 6). Compounds indicated by hatching are preferred.

20

Figure 1: Add-on treatment with pipamperon after treatment with citalopram .

Figure 2: HDRS-17 change from baseline: combo treatment pipamperon as add-on - citalopram vs SNRI (duloxetine) in Major Depression.

25

Figure 3: Remission rates (HDRS-17 ≤ 7): combo treatment pipamperon as add-on - citalopram vs SNRI (venlafaxine) vs SSRIs vs placebo in Major Depression.

30

Figure 4: Fore-going treatment during 1-5 days with pipamperon followed with the combination treatment of pipamperon and citalopram.

Figure 5: HDRS-17 change from baseline: combo treatment pipamperon - citalopram with a fore-going treatment of 4 days with pipamperon vs SNRI (duloxetine) in Major Depression.

35

Figure 6: Remission rates (HDRS-17 ≤ 7): combo pipamperon - citalopram with a fore-going treatment of 4 days with pipamperon vs SNRI (venlafaxine) in Major Depression.

5 **Figure 7:** Fore-going treatment during 6-8 days with pipamperon followed with the combination treatment of pipamperon and citalopram.

Figure 8: HDRS-17 change from baseline: combo treatment pipamperon - citalopram with a fore-going treatment of 7 days with pipamperon vs SNRI (duloxetine) in Major Depression.

10

Figure 9: Fore-going and add-on treatment with pipamperon in MDD.

Figure 10: HDRS-17 change from baseline: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SNRI duloxetine in Major Depression.

15

Figure 11: Remission rates (HDRS-17 ≤ 7): fore-going and add-on treatment with pipamperon and citalopram in comparison with the SNRI venlafaxine in Major Depression.

20 **Figure 12:** Y-BOCS total score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI fluvoxamine in OCD.

Figure 13: Y-BOCS obsession score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI fluvoxamine in OCD.

25 **Figure 14:** Y-BOCS compulsion score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI fluvoxamine in OCD.

Figure 15: CGI-severity score: fore-going and add-on treatment with pipamperon and citalopram in comparison with the SSRI in panic disorder.

Table 1

	D1	D2	D3	SHT _{1A}	SHT _{1B}	SHT _{1D}	SHT _{1E}	SHT _{1F}	SHT _{2A}	SHT _{2B}	SHT _{2C}	SHT _{6rat}	SHT _{7rat}	Alpha1A	Alpha2A	Alpha2B	Alpha2C	Beta1	Beta2	H1
ORG5222	8.9	8.9	8.9	8.9	8.9	8.9	7.8	0	8.9	8.9	8.9	8.9	8.9	8.9	8.9	8.9	7.8	<6	<6	<6
Zolopine	0	8.9	8.9	6.7	7.8	7.8	6.7	0	8.9	0	0	0	0	0	6.7	8.9	6.7	<6	<6	8.9
Fluparoxan	0	<6	<6	6.7	<6	<6	0	0	8.9	0	<6	0	0	6.7	8.9	8.9	8.9	0	0	0
Olanzapine	7.8	7.8	7.8	<6	6.7	6.7	<6	6.7	8.9	8.9	8.9	7.8	6.7	7.8	6.7	6.7	6.7	<6	<6	<6
Clozapine	7.8	6.7	6.7	6.7	6.7	6.7	6.7	6.7	8.9	8.9	7.8	7.8	7.8	8.9	7.8	7.8	7.8	<6	<6	<6
S16924	0	7.8	7.8	8.9	0	0	0	0	8.9	8.9	7.8	7.8	7.8	8.9	6.7	7.8	6.7	<6	<6	0
S18327	7.8	7.8	6.7	7.8	0	0	0	0	8.9	0	6.7	0	0	8.9	6.7	0	0	0	0	0
Amperozide	6.7	6.7	6.7	<6	0	0	0	0	8.9	0	<6	0	0	7.8	<6	0	0	0	0	0
GGR218231	<6	7.8	8.9	6.7	<6	<6	0	0	8.9	<6	<6	0	0	<6	<6	0	0	0	0	0
Sertindole	7.8	8.9	8.9	6.7	7.8	7.8	6.7	6.7	8.9	0	8.9	0	0	8.9	6.7	6.7	6.7	<6	<6	6.7
MDL100,907	6.7	<6	<6	<6	0	0	0	0	8.9	<6	<6	0	0	<6	<6	0	0	0	0	0
Haloperidol	8.9	8.9	8.9	<6	6.7	<6	<6	<6	8.9	<6	<6	<6	6.7	8.9	<6	6.7	<6	<6	<6	6.7
Tiospirone	7.8	8.9	8.9	8.9	0	0	0	0	8.9	0	8.9	0	0	8.9	6.7	0	0	0	0	0
Raclopride	<6	8.9	8.9	<6	0	0	0	0	8.9	0	<6	0	0	<6	<6	0	0	0	0	0
Fluspirilene	0	<6	8.9	7.8	<6	<6	<6	0	8.9	0	0	0	0	0	6.7	7.8	7.8	6.7	6.7	7.8
Ocaperidone	7.8	<6	8.9	7.8	0	0	0	0	8.9	0	7.8	0	0	8.9	0	0	0	0	0	0
Risperidone	7.8	8.9	7.8	6.7	8.9	6.7	<6	<6	8.9	0	7.8	0	0	8.9	7.8	8.9	8.9	<6	<6	7.8
S33084	6.7	7.8	8.9	<6	6.7	6.7	0	0	8.9	6.7	7.8	0	0	6.7	<6	0	0	0	0	0
L741626	6.7	8.9	7.8	<6	<6	<6	0	0	8.9	6.7	<6	0	0	6.7	<6	0	0	0	0	0
Seroquel	6.7	6.7	6.7	6.7	<6	<6	<6	<6	8.9	6.7	6.7	0	0	7.8	<6	7.8	6.7	<6	<6	8.9
Yohimbine	0	6.7	<6	7.8	6.7	7.8	0	0	8.9	0	<6	0	0	6.7	8.9	8.9	8.9	<6	<6	0
Ziprasidone	8.9	8.9	7.8	8.9	8.9	8.9	6.7	0	8.9	8.9	8.9	7.8	8.9	8.9	6.7	7.8	7.8	<6	<6	7.8
Pipamperon	0	6.7	6.7	<6	6.7	6.7	<6	<6	8.9	8.9	0	0	0	0	6.7	7.8	6.7	<6	<6	<6

Table 2

ACUTE PHASE**										EXTENSION PHASE***										FOLLOW-UP PHASE									
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
TREATMENT GROUP										TREATMENT GROUP										TREATMENT GROUP									
Group 1: Active D7										Group 1: Active D7										Group 1: Active D7									
Group 2: Active D4										Group 2: Active D4										Group 2: Active D4									
Group 3: Active D1										Group 3: Active D1										Group 3: Active D1									
Group 4: Active D0										Group 4: Active D0										Group 4: Active D0									
Group 5: Active D-1										Group 5: Active D-1										Group 5: Active D-1									
Group 6: Active D-2										Group 6: Active D-2										Group 6: Active D-2									
Group 7: Active D-3										Group 7: Active D-3										Group 7: Active D-3									
Group 8: Active D-4										Group 8: Active D-4										Group 8: Active D-4									
Group 9: Active D-5										Group 9: Active D-5										Group 9: Active D-5									
Group 10: Active D-6										Group 10: Active D-6										Group 10: Active D-6									
Group 11: Active D-7										Group 11: Active D-7										Group 11: Active D-7									
Group 12: Active D-8										Group 12: Active D-8										Group 12: Active D-8									
Group 13: Active D-9										Group 13: Active D-9										Group 13: Active D-9									
Group 14: Active D-10										Group 14: Active D-10										Group 14: Active D-10									
Group 15: Active D-11										Group 15: Active D-11										Group 15: Active D-11									
Group 16: Active D-12										Group 16: Active D-12										Group 16: Active D-12									
Group 17: Active D-13										Group 17: Active D-13										Group 17: Active D-13									
Group 18: Active D-14										Group 18: Active D-14										Group 18: Active D-14									
Group 19: Active D-15										Group 19: Active D-15										Group 19: Active D-15									
Group 20: Active D-16										Group 20: Active D-16										Group 20: Active D-16									
Group 21: Active D-17										Group 21: Active D-17										Group 21: Active D-17									
Group 22: Active D-18										Group 22: Active D-18										Group 22: Active D-18									
Group 23: Active D-19										Group 23: Active D-19										Group 23: Active D-19									
Group 24: Active D-20										Group 24: Active D-20										Group 24: Active D-20									
Group 25: Active D-21										Group 25: Active D-21										Group 25: Active D-21									
Group 26: Active D-22										Group 26: Active D-22										Group 26: Active D-22									
Group 27: Active D-23										Group 27: Active D-23										Group 27: Active D-23									
Group 28: Active D-24										Group 28: Active D-24										Group 28: Active D-24									
Group 29: Active D-25										Group 29: Active D-25										Group 29: Active D-25									
Group 30: Active D-26										Group 30: Active D-26										Group 30: Active D-26									
Group 31: Active D-27										Group 31: Active D-27										Group 31: Active D-27									
Group 32: Active D-28										Group 32: Active D-28										Group 32: Active D-28									
Group 33: Active D-29										Group 33: Active D-29										Group 33: Active D-29									
Group 34: Active D-30										Group 34: Active D-30										Group 34: Active D-30									
Group 35: Active D-31										Group 35: Active D-31										Group 35: Active D-31									
Group 36: Active D-32										Group 36: Active D-32										Group 36: Active D-32									
Group 37: Active D-33										Group 37: Active D-33										Group 37: Active D-33									
Group 38: Active D-34										Group 38: Active D-34										Group 38: Active D-34									
Group 39: Active D-35										Group 39: Active D-35										Group 39: Active D-35									
Group 40: Active D-36										Group 40: Active D-36										Group 40: Active D-36									
Group 41: Active D-37										Group 41: Active D-37										Group 41: Active D-37									
Group 42: Active D-38										Group 42: Active D-38										Group 42: Active D-38									
Group 43: Active D-39										Group 43: Active D-39										Group 43: Active D-39									
Group 44: Active D-40										Group 44: Active D-40										Group 44: Active D-40									
Group 45: Active D-41										Group 45: Active D-41										Group 45: Active D-41									
Group 46: Active D-42										Group 46: Active D-42										Group 46: Active D-42									
Group 47: Active D-43										Group 47: Active D-43										Group 47: Active D-43									
Group 48: Active D-44										Group 48: Active D-44										Group 48: Active D-44									
Group 49: Active D-45										Group 49: Active D-45										Group 49: Active D-45									
Group 50: Active D-46										Group 50: Active D-46										Group 50: Active D-46									
Group 51: Active D-47										Group 51: Active D-47										Group 51: Active D-47									
Group 52: Active D-48										Group 52: Active D-48										Group 52: Active D-48									
Group 53: Active D-49										Group 53: Active D-49										Group 53: Active D-49									
Group 54: Active D-50										Group 54: Active D-50										Group 54: Active D-50									
Group 55: Active D-51										Group 55: Active D-51										Group 55: Active D-51									
Group 56: Active D-52										Group 56: Active D-52										Group 56: Active D-52									
Group 57: Active D-53										Group 57: Active D-53										Group 57: Active D-53									
Group 58: Active D-54										Group 58: Active D-54										Group 58: Active D-54									
Group 59: Active D-55										Group 59: Active D-55										Group 59: Active D-55									
Group 60: Active D-56										Group 60: Active D-56										Group 60: Active D-56									
Group 61: Active D-57										Group 61: Active D-57										Group 61: Active D-57									
Group 62: Active D-58										Group 62: Active D-58										Group 62: Active D-58									
Group 63: Active D-59										Group 63: Active D-59										Group 63: Active D-59									
Group 64: Active D-60										Group 64: Active D-60										Group 64: Active D-60									
Group 65: Active D-61										Group 65: Active D-61										Group 65: Active D-61									
Group 66: Active D-62										Group 66: Active D-62										Group 66: Active D-62									
Group 67: Active D-63										Group 67: Active D-63										Group 67: Active D-63									
Group 68: Active D-64										Group 68: Active D-64										Group 68: Active D-64									
Group 69: Active D-65										Group 69: Active D-65										Group 69: Active D-65									
Group 70: Active D-66										Group 70: Active D-66										Group 70: Active D-66									
Group 71: Active D-67										Group 71: Active D-67										Group 71: Active D-67									
Group 72: Active D-68										Group 72: Active D-68										Group 72: Active D-68									
Group 73: Active D-69										Group 73: Active D-69										Group 73: Active D-69									
Group 74: Active D-70										Group 74: Active D-70										Group 74: Active D-70									
Group 75: Active D-71										Group 75: Active D-71										Group 75: Active D-71									
Group 76: Active D-72										Group 76: Active D-72										Group 76: Active D-72									
Group 77: Active D-73										Group 77: Active D-73										Group 77: Active D-73									
Group 78: Active D-74										Group 78: Active D-74										Group 78: Active D-74									
Group 79: Active D-75										Group 79: Active D-75										Group 79: Active D-75									
Group 80: Active D-76										Group 80: Active D-76										Group 80: Active D-76									
Group 81: Active D-77										Group 81: Active D-77										Group 81: Active D-77									
Group 82: Active D-78										Group 82: Active D-78										Group 82: Active D-78									
Group 83: Active D-79										Group 83: Active D-79										Group 83: Active D-79									
Group 84: Active D-80										Group 84: Active D-80										Group 84: Active D-80									
Group 85: Active D-81										Group 85: Active D-81										Group 85: Active D-81									
Group 86: Active D-82										Group 86: Active D-82										Group 86: Active D-82									
Group 87: Active D-83										Group 87: Active D-83										Group 87: Active D-83									
Group 88: Active D-84										Group 88: Active D-84										Group 88: Active D-84									
Group 89: Active D-85										Group 89: Active D-85										Group 89: Active D-85									
Group 90: Active D-86										Group 90: Active D-86										Group 90: Active D-86									
Group 91: Active D-87										Group 91: Active D-87										Group 91: Active D-87									
Group 92: Active D-88										Group 92: Active D-88										Group 92: Active D-88									
Group 93: Active D-89										Group 93: Active D-89										Group 93: Active D-89									
Group 94: Active D-90										Group 94: Active D-90										Group 94: Active D-90									
Group 95: Active D-91										Group 95: Active D-91										Group 95: Active D-91									
Group 96: Active D-92										Group 96: Active D-92										Group 96: Active D-92									
Group 97: Active D-93										Group 97: Active D-93										Group 97: Active D-93									
Group 98: Active D-94										Group 98: Active D-94										Group 98: Active D-94									
Group 99: Active D-95										Group 99: Active D-95										Group 99: Active D-95									
Group 100: Active D-96										Group 100: Active D-96										Group 100: Active D-96									
Group 101: Active D-97										Group 101: Active D-97										Group 101: Active D-97									
Group 102: Active D-98										Group 102: Active D-98										Group 102: Active D-98									
Group 103: Active D-99										Group 103: Active D-99										Group 103: Active D-99									
Group 104: Active D-100										Group 104: Active D-100										Group 104: Active D-100									
Group 105: Active D-101										Group 105: Active D-101										Group 105: Active D-101									
Group 106: Active D-102										Group 106: Active D-102										Group 106: Active D-102									
Group 107: Active D-103										Group 107: Active D-103										Group 107: Active D-103									
Group 108: Active D-104										Group 108: Active D-104										Group 108: Active D-104									
Group 109: Active D-105										Group 109: Active D-105										Group 109: Active D-105									
Group 110: Active D-106										Group 110: Active D-106										Group 110: Active D-106									
Group 111: Active D-107										Group 111: Active D-107										Group 111: Active D-107									
Group 112: Active D-108										Group 112: Active D-108										Group 112: Active D-108									
Group 113: Active D-109										Group 113: Active D-109										Group 113: Active D-109									
Group 114: Active D-110										Group 114: Active D-110										Group 114: Active D-110									
Group 115: Active D-111										Group 115: Active D-111										Group 115: Active D-111									
Group 116: Active D-112										Group 116: Active D-112										Group 116: Active D-112									
Group 117: Active D-113										Group 117: Active D-113										Group 117: Active D-113									
Group 118: Active D-114										Group 118: Active D-114										Group 118: Active D-114									
Group 119: Active D-115										Group 119: Active D-115										Group 119: Active D-115									
Group 120: Active D-116										Group 120: Active D-116										Group 120: Active D-116									
Group 121: Active D-117										Group 121: Active D-117										Group 121: Active D-117									
Group 122: Active D-118										Group 122: Active D-118										Group 122: Active D-118									
Group 123: Active D-119										Group 123: Active D-119										Group 123: Active D-119									
Group 124: Active D-120										Group 124: Active D-120										Group 124: Active D-120									
Group 125: Active D-121										Group 125: Active D-121										Group 125: Active D-121									
Group 126: Active D-122										Group 126: Active D-122										Group 126: Active D-122									
Group 127: Active D-123										Group 127: Active D-123										Group 127: Active D-123									
Group 128: Active D-124										Group 128: Active D-124										Group 128: Active D-124									
Group 129: Active D-125										Group 129: Active D-125										Group 129: Active D-125									
Group 130: Active D-126										Group 130: Active D-126										Group 130: Active D-126									
Group 131: Active D-127										Group 131: Active D-127										Group 131: Active D-127									
Group 132: Active D-128										Group 132: Active D-128										Group 132: Active D-128									
Group 133: Active D-129										Group 133: Active D-129										Group 133: Active D-129									
Group 134: Active D-130										Group 134: Active D-130										Group 134: Active D-130									
Group 135: Active D-131										Group 135: Active D-131										Group 135: Active D-131									
Group 136: Active D-132										Group 136: Active D-132										Group 136: Active D-132									
Group 137: Active D-133										Group 137: Active D-133										Group 137: Active D-133									
Group 138: Active D-134										Group 138: Active D-134										Group 138: Active D-134									
Group 139: Active D-135										Group 139: Active D-135										Group 139: Active D-135									
Group 140: Active D-136										Group 140: Active D-136										Group 140: Active D-136									
Group 141: Active D-137										Group 141: Active D-137										Group 141: Active D-137									
Group 142: Active D-138										Group 142: Active D-138										Group 142: Active D-138									
Group 143: Active D-139										Group 143: Active D-139										Group 143: Active D-139									
Group 144: Active D-140										Group 144: Active D-140										Group 144: Active D-140									
Group 145: Active D-141										Group 145: Active D-141										Group 145: Active D-141									
Group 146: Active D-142										Group 146: Active D-142										Group 146: Active D-142									
Group 147: Active D-143										Group 147: Active D-143										Group 147: Active D-143									
Group 148: Active D-144										Group 148: Active D-144										Group 148: Active D-144									
Group 149: Active D-145										Group 149: Active D-145										Group 149: Active D-145									
Group 150: Active D-146										Group 150: Active D-146										Group 150: Active D-146									
Group 151: Active D-147										Group 151: Active D-147										Group 151: Active D-147									
Group 152: Active D-148										Group 152: Active D-148										Group 152: Active D-148									
Group 153: Active D-149										Group 153: Active D-149										Group 153: Active D-149									
Group 154: Active D-150										Group 154: Active D-150										Group 154: Active D-150									
Group 155: Active D-151										Group 155: Active D-151										Group 155: Active D-151									
Group 156: Active D-152										Group 156: Active D-152										Group 156: Active D-152									
Group 157: Active D-153										Group 157: Active D-153										Group 157: Active D-153									
Group 158: Active D-154										Group 158: Active D-154										Group 158: Active D-154									
Group 159: Active D-155										Group 159: Active D-155										Group 159: Active D-155									
Group 160: Active D-156										Group 160: Active D-156										Group 160: Active D-156									
Group 161: Active D-157										Group 161: Active D-157										Group 161: Active D-157									
Group 162: Active D-158										Group 162: Active D-158										Group 162: Active D-158									
Group 163: Active D-159										Group 163: Active D-159										Group 163: Active D-159									
Group 164: Active D-160										Group 164: Active D-160										Group 164: Active D-160									
Group 165: Active D-161										Group 165: Active D-161										Group 165: Active D-161									
Group 166: Active D-162										Group 166: Active D-162										Group 166: Active D-162									
Group 167: Active D-163										Group 167: Active D-163										Group 167: Active D-163									
Group 168: Active D-164										Group 168: Active D-164										Group 168: Active D-164									
Group 169: Active D-165										Group 169: Active D-165										Group 169: Active D-165									
Group 170: Active D-166										Group 170: Active D-166										Group 170: Active D-166									
Group 171: Active D-167										Group 171: Active D-167										Group 171: Active D-167									
Group 172: Active D-168										Group 172: Active D-168										Group 172: Active D-168									
Group 173: Active D-169										Group 173: Active D-169										Group 173: Active D-169									
Group 174: Active D-170										Group 174: Active D-170										Group 174: Active D-170									
Group 175: Active D-171										Group 175: Active D-171										Group 175: Active D-171									
Group 176: Active D-172										Group 176: Active D-172										Group 176: Active D-172									
Group 177: Active D-173										Group 177: Active D-173										Group 177: Active D-173									
Group 178: Active D-174										Group 178: Active D-174										Group 178: Active D-174									
Group 179: Active D-175										Group 179: Active D-175										Group 179: Active D-175									
Group 180: Active D-176										Group 180: Active D-176										Group 180: Active D-176									
Group 181: Active D-177										Group 181: Active D-177										Group 181: Active D-177									
Group 182: Active D-178										Group 182: Active D-178										Group 182: Active D-178									
Group 183: Active D-179										Group 183: Active D-179										Group 183: Active D-179									
Group 184: Active D-180										Group 184: Active D-180										Group 184: Active D-180									
Group 185: Active D-181										Group 185: Active D-181										Group 185: Active D-181									
Group 186: Active D-182										Group 186: Active D-182										Group 186: Active D-182									
Group 187: Active D-183										Group 187: Active D-183										Group 187: Active D-183									
Group 188: Active D-184										Group 188: Active D-184										Group 188: Active D-184									
Group 189: Active D-185										Group 189: Active D-185										Group 189: Active D-185									
Group 190: Active D-186										Group 190: Active D-186										Group 190: Active D-186									
Group 191: Active D-187										Group 191: Active D-187																			

Table 3

FOREGOING PIPAMPERON - CITALOPRAM TREATMENT IN MAJOR DEPRESSIVE DISORDER									
A PLACEBO ACTIVE AND PERIODIC CONTROLLED									
CLINICAL TRIAL									
ACUTE PHASE**					EXTENSION PHASE***				
FOLLOW-UP PHASE									
DAY	WEEK	MONTH	YEAR	GROUP	DAY	WEEK	MONTH	YEAR	GROUP
1	1	1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2	2	2
3	3	3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4	4	4
5	5	5	5	5	5	5	5	5	5
6	6	6	6	6	6	6	6	6	6
7	7	7	7	7	7	7	7	7	7
8	8	8	8	8	8	8	8	8	8
9	9	9	9	9	9	9	9	9	9
10	10	10	10	10	10	10	10	10	10
11	11	11	11	11	11	11	11	11	11
12	12	12	12	12	12	12	12	12	12
13	13	13	13	13	13	13	13	13	13
14	14	14	14	14	14	14	14	14	14
15	15	15	15	15	15	15	15	15	15
16	16	16	16	16	16	16	16	16	16
17	17	17	17	17	17	17	17	17	17
18	18	18	18	18	18	18	18	18	18
19	19	19	19	19	19	19	19	19	19
20	20	20	20	20	20	20	20	20	20
21	21	21	21	21	21	21	21	21	21
22	22	22	22	22	22	22	22	22	22
23	23	23	23	23	23	23	23	23	23
24	24	24	24	24	24	24	24	24	24
25	25	25	25	25	25	25	25	25	25
26	26	26	26	26	26	26	26	26	26
27	27	27	27	27	27	27	27	27	27
28	28	28	28	28	28	28	28	28	28
29	29	29	29	29	29	29	29	29	29
30	30	30	30	30	30	30	30	30	30
31	31	31	31	31	31	31	31	31	31
32	32	32	32	32	32	32	32	32	32
33	33	33	33	33	33	33	33	33	33
34	34	34	34	34	34	34	34	34	34
35	35	35	35	35	35	35	35	35	35
36	36	36	36	36	36	36	36	36	36
37	37	37	37	37	37	37	37	37	37
38	38	38	38	38	38	38	38	38	38
39	39	39	39	39	39	39	39	39	39
40	40	40	40	40	40	40	40	40	40
41	41	41	41	41	41	41	41	41	41
42	42	42	42	42	42	42	42	42	42
43	43	43	43	43	43	43	43	43	43
44	44	44	44	44	44	44	44	44	44
45	45	45	45	45	45	45	45	45	45
46	46	46	46	46	46	46	46	46	46
47	47	47	47	47	47	47	47	47	47
48	48	48	48	48	48	48	48	48	48
49	49	49	49	49	49	49	49	49	49
50	50	50	50	50	50	50	50	50	50
51	51	51	51	51	51	51	51	51	51
52	52	52	52	52	52	52	52	52	52
53	53	53	53	53	53	53	53	53	53
54	54	54	54	54	54	54	54	54	54
55	55	55	55	55	55	55	55	55	55
56	56	56	56	56	56	56	56	56	56
57	57	57	57	57	57	57	57	57	57
58	58	58	58	58	58	58	58	58	58
59	59	59	59	59	59	59	59	59	59
60	60	60	60	60	60	60	60	60	60
61	61	61	61	61	61	61	61	61	61
62	62	62	62	62	62	62	62	62	62
63	63	63	63	63	63	63	63	63	63
64	64	64	64	64	64	64	64	64	64
65	65	65	65	65	65	65	65	65	65
66	66	66	66	66	66	66	66	66	66
67	67	67	67	67	67	67	67	67	67
68	68	68	68	68	68	68	68	68	68
69	69	69	69	69	69	69	69	69	69
70	70	70	70	70	70	70	70	70	70
71	71	71	71	71	71	71	71	71	71
72	72	72	72	72	72	72	72	72	72
73	73	73	73	73	73	73	73	73	73
74	74	74	74	74	74	74	74	74	74
75	75	75	75	75	75	75	75	75	75
76	76	76	76	76	76	76	76	76	76
77	77	77	77	77	77	77	77	77	77
78	78	78	78	78	78	78	78	78	78
79	79	79	79	79	79	79	79	79	79
80	80	80	80	80	80	80	80	80	80
81	81	81	81	81	81	81	81	81	81
82	82	82	82	82	82	82	82	82	82
83	83	83	83	83	83	83	83	83	83
84	84	84	84	84	84	84	84	84	84
85	85	85	85	85	85	85	85	85	85
86	86	86	86	86	86	86	86	86	86
87	87	87	87	87	87	87	87	87	87
88	88	88	88	88	88	88	88	88	88
89	89	89	89	89	89	89	89	89	89
90	90	90	90	90	90	90	90	90	90
91	91	91	91	91	91	91	91	91	91
92	92	92	92	92	92	92	92	92	92
93	93	93	93	93	93	93	93	93	93
94	94	94	94	94	94	94	94	94	94
95	95	95	95	95	95	95	95	95	95
96	96	96	96	96	96	96	96	96	96
97	97	97	97	97	97	97	97	97	97
98	98	98	98	98	98	98	98	98	98
99	99	99	99	99	99	99	99	99	99
100	100	100	100	100	100	100	100	100	100

*Neuronal E-Clinical Trial = Vesalius Expert Development for this Trial which includes the bottom-up measurement of:

- . In- and exclusion criteria
- . Functional status evaluation
- . Medical history
- . (Pre-)treatment signs & symptoms
- . DSM-IV rules for diagnosis & efficacy
- . Rating Scales: HDRS-21, MADRS, HAM-A
- . Medical resource utilisation
- . Pre-treatment concomitant medication
- . Drug administration
- . (Serious) Adverse events
- . Admission to the acute and extension phase of treatment
- . Right flow of the trial

**Entering Acute Phase: only NON-placebo responders as defined by the DSM-IV criteria of efficacy

***Entering Extension Phase: only remitters as defined by the DSM-IV criteria of efficacy

****Q-LES-Q: Quality of Life, Enjoyment and Satisfaction Questionnaire

179

Table 4

5

DAY	minus D7	D0	=>D4
TREATMENTGROUP			
Placebo (PLC)	PLC+PLC	2x(PLC+PLC)	2x(PLC+PC)
PIP - Active / Day 4	PLC+PLC	2x(PLC+PIP (4mg))/d	2x(CIT (10mg)+PIP (4mg))/d
PIP - Active / Day 0	PLC+PLC	2x(CIT (10mg)+PIP (4mg))/d	2x(CIT (10mg)+PIP (4mg))/d
PLC - Active / Day 0	PLC+PLC	2x(CIT (10mg)+PLC)/d	2x(CIT (10mg)+PLC)/d

10

החברות הן:

[illegible]

[illegible]

GLOSSARY	
5-HT	= serotonin
5-HT ₁	= serotonin 1 receptor
5-HT _{1A}	= serotonin 1A receptor
5-HT _{1B}	= serotonin 1B receptor
5-HT _{2A/D4}	= serotonin 2A and dopamine D4 receptor
5-HT _{2B}	= serotonin 2B receptor
5-HT _{2C}	= serotonin 2C receptor
5-HT ₃	= serotonin 3 receptor
5-HT ₆	= serotonin 6 receptor
AMPA	= α -l-glutamate-3-hydroxy-5-methyl-4-isoxazole propionate
CB ₁	= cannabinoid receptor 1
CINODs	= COX-inhibiting nitric oxide donors
COX	= cyclooxygenase
COX-2	= cyclooxygenase 2
CRF-1	= Corticotropin-Releasing Factor Receptor 1
D ₁	= Dopamine 1
D ₂	= Dopamine 2
D ₃	= Dopamine 3
DA	= Dopamine
ERK	= extracellular signal-related kinase
GABA	= gamma-aminobutyric acid
GABA-A	= gamma-aminobutyric acid A receptor
GABA-B	= gamma-aminobutyric acid B receptor
GPCR	= G-Protein-Coupled Receptor
GR	= glucocorticoid receptor
H ₃	= histamine H ₃ -receptor
MAO	= mono-amine oxidase
MAO-A	= mono-amine oxidase A
MAO-B	= mono-amine oxidase B
MCH	= melanin-concentrating hormone
MCH ₁	= Melanin-concentrating hormone
MolR	= melanopsin melanopsin receptor
MT	= melanin receptor
NARI	= selective nor-adrenaline re-uptake inhibitor
NaSSA	= noradrenergic/specific serotonergic antidepressant
NDRI	= selective nor-adrenaline and dopamine re-uptake inhibitor
NGF	= Nerve Growth Factor
NGF	= Nerve growth factor
NK ₁	= neurokinin 1 receptor
NK ₂	= neurokinin 2 receptor
NK ₃	= neurokinin 3 receptor
NMDA	= N-Methyl-D-aspartate
NSAID	= Non-steroidal anti-inflammatory drugs
PDE4	= phosphodiesterase-4
RIMA	= reversible inhibitor of mono-amine oxidase A
SCT-11	= G protein-coupled receptor
SDA	= Serotonin/Dopamine Antagonist
SDRI	= selective serotonin and dopamine reuptake inhibitor
SNRI	= selective serotonin, nor-adrenaline and dopamine reuptake inhibitor
SNRI	= selective serotonin and nor-adrenaline reuptake inhibitor
SSRI	= selective serotonin reuptake inhibitor
VIB	= vasopressin 1B receptor

188

PHARMAC. GROUP (see overview bar/row)	nr. PH. PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Monoaminergic Transmitter Systems	1	5-HT _{1A} agonist/enhancer	Depression/Anxiety	1025551	100 mg/day	Novartis
	2	5-HT _{1A} antagonist	Depression/Anxiety	1025552	100 mg/day	Novartis
	3	5-HT _{1A} agonist	Anxiety	1025553	100 mg/day	Novartis
	4	5-HT _{1A} agonist	Depression/Anxiety	Bupropion	300 mg/day	Bristol-Myers Squibb
	5	5-HT _{1A} agonist	Depression	1025554	100 mg/day	Novartis
	6	5-HT _{1A} agonist	Alzheimer's Disease	1025555	100 mg/day	Novartis
	7	5-HT _{1A} agonist	Depression/Anxiety	1025556	100 mg/day	Novartis
	8	5-HT _{1A} agonist	Anxiety	1025557	100 mg/day	Novartis
	9	5-HT _{1A} agonist	Depression	1025558	100 mg/day	Novartis
	10	5-HT _{1A} agonist	Depression/Anxiety	1025559	100 mg/day	Novartis
	11	5-HT _{1A} agonist	Depression/Anxiety	1025560	100 mg/day	Novartis
	12	5-HT _{1A} agonist	Depression/Anxiety	1025561	100 mg/day	Novartis
	13	5-HT _{1A} agonist	Depression/Anxiety	1025562	100 mg/day	Novartis
	14	5-HT _{1A} agonist	Depression/Anxiety	1025563	100 mg/day	Novartis
	15	5-HT _{1A} agonist	Depression/Anxiety	1025564	100 mg/day	Novartis
Monoaminergic Transmitter Systems	16	5-HT _{1B} agonist	Depression/Anxiety	1025565	100 mg/day	Novartis
	17	5-HT _{1B} agonist	Depression/Anxiety	1025566	100 mg/day	Novartis
	18	5-HT _{1B} agonist	Depression/Anxiety	1025567	100 mg/day	Novartis
	19	5-HT _{1B} agonist	Depression/Anxiety	1025568	100 mg/day	Novartis
	20	5-HT _{1B} agonist	Depression/Anxiety	1025569	100 mg/day	Novartis
	21	5-HT _{1B} agonist	Depression/Anxiety	1025570	100 mg/day	Novartis
	22	5-HT _{1B} agonist	Depression/Anxiety	1025571	100 mg/day	Novartis
	23	5-HT _{1B} agonist	Depression/Anxiety	1025572	100 mg/day	Novartis
	24	5-HT _{1B} agonist	Depression/Anxiety	1025573	100 mg/day	Novartis
	25	5-HT _{1B} agonist	Depression/Anxiety	1025574	100 mg/day	Novartis
	26	5-HT _{1B} agonist	Depression/Anxiety	1025575	100 mg/day	Novartis
	27	5-HT _{1B} agonist	Depression/Anxiety	1025576	100 mg/day	Novartis
	28	5-HT _{1B} agonist	Depression/Anxiety	1025577	100 mg/day	Novartis
	29	5-HT _{1B} agonist	Depression/Anxiety	1025578	100 mg/day	Novartis
	30	5-HT _{1B} agonist	Depression/Anxiety	1025579	100 mg/day	Novartis
Monoaminergic Transmitter Systems	31	5-HT _{1C} agonist	Depression/Anxiety	1025580	100 mg/day	Novartis
	32	5-HT _{1C} agonist	Depression/Anxiety	1025581	100 mg/day	Novartis
	33	5-HT _{1C} agonist	Depression/Anxiety	1025582	100 mg/day	Novartis
	34	5-HT _{1C} agonist	Depression/Anxiety	1025583	100 mg/day	Novartis
	35	5-HT _{1C} agonist	Depression/Anxiety	1025584	100 mg/day	Novartis
	36	5-HT _{1C} agonist	Depression/Anxiety	1025585	100 mg/day	Novartis
	37	5-HT _{1C} agonist	Depression/Anxiety	1025586	100 mg/day	Novartis
Monoaminergic Transmitter Systems	38	5-HT _{1D} agonist	Depression/Anxiety	1025587	100 mg/day	Novartis
	39	5-HT _{1D} agonist	Depression/Anxiety	1025588	100 mg/day	Novartis
	40	5-HT _{1D} agonist	Depression/Anxiety	1025589	100 mg/day	Novartis
	41	5-HT _{1D} agonist	Depression/Anxiety	1025590	100 mg/day	Novartis
	42	5-HT _{1D} agonist	Depression/Anxiety	1025591	100 mg/day	Novartis
	43	5-HT _{1D} agonist	Depression/Anxiety	1025592	100 mg/day	Novartis
	44	5-HT _{1D} agonist	Depression/Anxiety	1025593	100 mg/day	Novartis
Monoaminergic Transmitter Systems	45	5-HT _{1E} agonist	Depression/Anxiety	1025594	100 mg/day	Novartis
	46	5-HT _{1E} agonist	Depression/Anxiety	1025595	100 mg/day	Novartis
	47	5-HT _{1E} agonist	Depression/Anxiety	1025596	100 mg/day	Novartis
	48	5-HT _{1E} agonist	Depression/Anxiety	1025597	100 mg/day	Novartis
	49	5-HT _{1E} agonist	Depression/Anxiety	1025598	100 mg/day	Novartis
	50	5-HT _{1E} agonist	Depression/Anxiety	1025599	100 mg/day	Novartis
	51	5-HT _{1E} agonist	Depression/Anxiety	1025600	100 mg/day	Novartis
	52	5-HT _{1E} agonist	Depression/Anxiety	1025601	100 mg/day	Novartis
	53	5-HT _{1E} agonist	Depression/Anxiety	1025602	100 mg/day	Novartis
	54	5-HT _{1E} agonist	Depression/Anxiety	1025603	100 mg/day	Novartis
	55	5-HT _{1E} agonist	Depression/Anxiety	1025604	100 mg/day	Novartis
	56	5-HT _{1E} agonist	Depression/Anxiety	1025605	100 mg/day	Novartis
	57	5-HT _{1E} agonist	Depression/Anxiety	1025606	100 mg/day	Novartis
	58	5-HT _{1E} agonist	Depression/Anxiety	1025607	100 mg/day	Novartis
	59	5-HT _{1E} agonist	Depression/Anxiety	1025608	100 mg/day	Novartis
Monoaminergic Transmitter Systems	60	5-HT _{1F} agonist	Depression/Anxiety	1025609	100 mg/day	Novartis
	61	5-HT _{1F} agonist	Depression/Anxiety	1025610	100 mg/day	Novartis
	62	5-HT _{1F} agonist	Depression/Anxiety	1025611	100 mg/day	Novartis
	63	5-HT _{1F} agonist	Depression/Anxiety	1025612	100 mg/day	Novartis
	64	5-HT _{1F} agonist	Depression/Anxiety	1025613	100 mg/day	Novartis
	65	5-HT _{1F} agonist	Depression/Anxiety	1025614	100 mg/day	Novartis
	66	5-HT _{1F} agonist	Depression/Anxiety	1025615	100 mg/day	Novartis
	67	5-HT _{1F} agonist	Depression/Anxiety	1025616	100 mg/day	Novartis
	68	5-HT _{1F} agonist	Depression/Anxiety	1025617	100 mg/day	Novartis
	69	5-HT _{1F} agonist	Depression/Anxiety	1025618	100 mg/day	Novartis
	70	5-HT _{1F} agonist	Depression/Anxiety	1025619	100 mg/day	Novartis
	71	5-HT _{1F} agonist	Depression/Anxiety	1025620	100 mg/day	Novartis
	72	5-HT _{1F} agonist	Depression/Anxiety	1025621	100 mg/day	Novartis
	73	5-HT _{1F} agonist	Depression/Anxiety	1025622	100 mg/day	Novartis
	74	5-HT _{1F} agonist	Depression/Anxiety	1025623	100 mg/day	Novartis
Monoaminergic Transmitter Systems	75	5-HT _{1G} agonist	Depression/Anxiety	1025624	100 mg/day	Novartis
	76	5-HT _{1G} agonist	Depression/Anxiety	1025625	100 mg/day	Novartis
	77	5-HT _{1G} agonist	Depression/Anxiety	1025626	100 mg/day	Novartis
	78	5-HT _{1G} agonist	Depression/Anxiety	1025627	100 mg/day	Novartis
	79	5-HT _{1G} agonist	Depression/Anxiety	1025628	100 mg/day	Novartis
	80	5-HT _{1G} agonist	Depression/Anxiety	1025629	100 mg/day	Novartis
	81	5-HT _{1G} agonist	Depression/Anxiety	1025630	100 mg/day	Novartis
	82	5-HT _{1G} agonist	Depression/Anxiety	1025631	100 mg/day	Novartis
	83	5-HT _{1G} agonist	Depression/Anxiety	1025632	100 mg/day	Novartis
	84	5-HT _{1G} agonist	Depression/Anxiety	1025633	100 mg/day	Novartis
	85	5-HT _{1G} agonist	Depression/Anxiety	1025634	100 mg/day	Novartis
	86	5-HT _{1G} agonist	Depression/Anxiety	1025635	100 mg/day	Novartis
	87	5-HT _{1G} agonist	Depression/Anxiety	1025636	100 mg/day	Novartis
	88	5-HT _{1G} agonist	Depression/Anxiety	1025637	100 mg/day	Novartis
	89	5-HT _{1G} agonist	Depression/Anxiety	1025638	100 mg/day	Novartis
Monoaminergic Transmitter Systems	90	5-HT _{1H} agonist	Depression/Anxiety	1025639	100 mg/day	Novartis
	91	5-HT _{1H} agonist	Depression/Anxiety	1025640	100 mg/day	Novartis
	92	5-HT _{1H} agonist	Depression/Anxiety	1025641	100 mg/day	Novartis
	93	5-HT _{1H} agonist	Depression/Anxiety	1025642	100 mg/day	Novartis
	94	5-HT _{1H} agonist	Depression/Anxiety	1025643	100 mg/day	Novartis
	95	5-HT _{1H} agonist	Depression/Anxiety	1025644	100 mg/day	Novartis
	96	5-HT _{1H} agonist	Depression/Anxiety	1025645	100 mg/day	Novartis
	97	5-HT _{1H} agonist	Depression/Anxiety	1025646	100 mg/day	Novartis
	98	5-HT _{1H} agonist	Depression/Anxiety	1025647	100 mg/day	Novartis
	99	5-HT _{1H} agonist	Depression/Anxiety	1025648	100 mg/day	Novartis
	100	5-HT _{1H} agonist	Depression/Anxiety	1025649	100 mg/day	Novartis
	101	5-HT _{1H} agonist	Depression/Anxiety	1025650	100 mg/day	Novartis
	102	5-HT _{1H} agonist	Depression/Anxiety	1025651	100 mg/day	Novartis
	103	5-HT _{1H} agonist	Depression/Anxiety	1025652	100 mg/day	Novartis
	104	5-HT _{1H} agonist	Depression/Anxiety	1025653	100 mg/day	Novartis
Monoaminergic Transmitter Systems	105	5-HT _{1I} agonist	Depression/Anxiety	1025654	100 mg/day	Novartis
	106	5-HT _{1I} agonist	Depression/Anxiety	1025655	100 mg/day	Novartis
	107	5-HT _{1I} agonist	Depression/Anxiety	1025656	100 mg/day	Novartis
	108	5-HT _{1I} agonist	Depression/Anxiety	1025657	100 mg/day	Novartis
	109	5-HT _{1I} agonist	Depression/Anxiety	1025658	100 mg/day	Novartis
	110	5-HT _{1I} agonist	Depression/Anxiety	1025659	100 mg/day	Novartis
	111	5-HT _{1I} agonist	Depression/Anxiety	1025660	100 mg/day	Novartis
	112	5-HT _{1I} agonist	Depression/Anxiety	1025661	100 mg/day	Novartis
	113	5-HT _{1I} agonist	Depression/Anxiety	1025662	100 mg/day	Novartis
	114	5-HT _{1I} agonist	Depression/Anxiety	1025663	100 mg/day	Novartis
	115	5-HT _{1I} agonist	Depression/Anxiety	1025664	100 mg/day	Novartis
	116	5-HT _{1I} agonist	Depression/Anxiety	1025665	100 mg/day	Novartis
	117	5-HT _{1I} agonist	Depression/Anxiety	1025666	100 mg/day	Novartis
	118	5-HT _{1I} agonist	Depression/Anxiety	1025667	100 mg/day	Novartis
	119	5-HT _{1I} agonist	Depression/Anxiety	1025668	100 mg/day	Novartis
Monoaminergic Transmitter Systems	120	5-HT _{1J} agonist	Depression/Anxiety	1025669	100 mg/day	Novartis
	121	5-HT _{1J} agonist	Depression/Anxiety	1025670	100 mg/day	Novartis
	122	5-HT _{1J} agonist	Depression/Anxiety	1025671	100 mg/day	Novartis
	123	5-HT _{1J} agonist	Depression/Anxiety	1025672	100 mg/day	Novartis
	124	5-HT _{1J} agonist	Depression/Anxiety	1025673	100 mg/day	Novartis
	125	5-HT _{1J} agonist	Depression/Anxiety	1025674	100 mg/day	Novartis
	126	5-HT _{1J} agonist	Depression/Anxiety	1025675	100 mg/day	Novartis
	127	5-HT _{1J} agonist	Depression/Anxiety	1025676	100 mg/day	Novartis
	128	5-HT _{1J} agonist	Depression/Anxiety	1025677	100 mg/day	Novartis
	129	5-HT _{1J} agonist	Depression/Anxiety	1025678	100 mg/day	Novartis
	130	5-HT _{1J} agonist	Depression/Anxiety	1025679	100 mg/day	Novartis
	131	5-HT _{1J} agonist	Depression/Anxiety	1025680	100 mg/day	Novartis
	132	5-HT _{1J} agonist	Depression/Anxiety	1025681	100 mg/day	Novartis
	133	5-HT _{1J} agonist	Depression/Anxiety	1025682	100 mg/day	Novartis
	134	5-HT _{1J} agonist	Depression/Anxiety	1025683	100 mg/day	Novartis
Monoaminergic Transmitter Systems	135	5-HT _{1K} agonist	Depression/Anxiety	1025684	100 mg/day	Novartis
	136	5-HT _{1K} agonist	Depression/Anxiety	1025685	100 mg/day	Novartis
	137	5-HT _{1K} agonist	Depression/Anxiety	1025686	100 mg/day	Novartis
	138	5-HT _{1K} agonist	Depression/Anxiety	1025687	100 mg/day	Novartis
	139	5-HT _{1K} agonist	Depression/Anxiety	1025688	100 mg/day	Novartis
	140	5-HT _{1K} agonist	Depression/Anxiety	1025689	100 mg/day	Novartis
	141	5-HT _{1K} agonist	Depression/Anxiety	1025690	100 mg/day	Novartis
	142	5-HT _{1K} agonist	Depression/Anxiety	1025691	100 mg/day	Novartis
	143	5-HT _{1K} agonist	Depression/Anxiety	1025692	100 mg/day	Novartis
	144	5-HT _{1K} agonist	Depression/Anxiety	1025693	100 mg/day	Novartis
	145	5-HT _{1K} agonist	Depression/Anxiety	1025694	100 mg/day	Novartis
	146	5-HT _{1K} agonist	Depression/Anxiety	1025695	100 mg/day	Novartis
	147	5-HT _{1K} agonist	Depression/Anxiety	1025696	100 mg/day	Novartis
	148	5-HT _{1K} agonist	Depression/Anxiety	1025697	100 mg/day	Novartis
	149	5-HT _{1K} agonist	Depression/Anxiety	1025698	100 mg/day	Novartis
Monoaminergic Transmitter Systems	150	5-HT _{1L} agonist	Depression/Anxiety	1025699	100 mg/day	Novartis
	151	5-HT _{1L} agonist	Depression/Anxiety	1025700	100 mg/day	Novartis
	152	5-HT _{1L} agonist	Depression/Anxiety	1025701	100 mg/day	Novartis
	153	5-HT _{1L} agonist	Depression/Anxiety	1025702	100 mg/day	Novartis
	154	5-HT _{1L} agonist	Depression/Anxiety	1025703	100 mg/day	Novartis
	155	5-HT _{1L} agonist	Depression/Anxiety	1025704	100 mg/day	Novartis
	156	5-HT _{1L} agonist	Depression/Anxiety	1025705	100 mg/day	Novartis
	157	5-HT _{1L} agonist	Depression/Anxiety	1025706	100 mg/day	Novartis
	158	5-HT _{1L} agonist	Depression/Anxiety	1025707	100 mg/day	Novartis
	159	5-HT _{1L} agonist	Depression/Anxiety	1025708	100 mg/day	Novartis
	160	5-HT _{1L} agonist	Depression/Anxiety	1025709	100 mg/day	Novartis
	161	5-HT _{1L} agonist	Depression/Anxiety	1025710	100 mg/day	Novartis
	162	5-HT _{1L} agonist	Depression/Anxiety	1025711	100 mg/day	Novartis
	163	5-HT _{1L} agonist	Depression/Anxiety	1025712	100 mg/day	Novartis
	164	5-HT _{1L} agonist	Depression/Anxiety	1025713	100 mg/day	Novartis
Monoaminergic Transmitter Systems	165	5-HT _{1M} agonist	Depression/Anxiety	1025714	100 mg/day	Novartis
	166	5-HT _{1M} agonist	Depression/Anxiety	1025715	100 mg/day	Novartis
	167	5-HT _{1M} agonist	Depression/Anxiety	1025716	100 mg/day	Novartis
	168	5-HT _{1M} agonist	Depression/Anxiety	1025717	100 mg/day	Novartis
	169	5-HT _{1M} agonist	Depression/Anxiety	1025718	100 mg/day	Novartis
	170	5-HT _{1M} agonist	Depression/Anxiety	1025719	100 mg/day	Novartis
	171	5-HT _{1M} agonist	Depression/Anxiety	1025720	100 mg/day	Novartis
	172	5-HT _{1M} agonist	Depression/Anxiety	1025721	100 mg/day	Novartis
	173	5-HT _{1M} agonist	Depression/Anxiety	1025722	100 mg/day	Novartis
	174	5-HT _{1M} agonist	Depression/Anxiety	1025723	100 mg/day	Novartis
	175	5-HT _{1M} agonist	Depression/Anxiety	1025724	100 mg/day	Novartis
	176	5-HT _{1M} agonist	Depression/Anxiety	1025725	100 mg/day	Novartis
	177	5-HT _{1M} agonist	Depression/Anxiety	1025726	100 mg/day	Novartis
	178	5-HT _{1M} agonist	Depression/Anxiety	1025727	100 mg/day	Novartis
	179	5-HT _{1M} agonist	Depression/Anxiety	1025728	100 mg/day	Novartis
Monoaminergic Transmitter Systems	180	5-HT _{1N} agonist	Depression/Anxiety	1025729		

[illegible]

[illegible]

PHARMAC. GROUP (see overview hereunder)	nr. PH. PROF.	PHARMACOLOGICAL PROFILE	MAIN INDICATIONS	COMPOUND	DOSE RANGE	COMPANY
Monoamine Transporter System	92	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	93	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	94	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	95	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	96	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	97	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	98	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	99	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	100	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	101	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	102	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	103	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	104	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	105	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	106	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	107	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	108	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	109	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	110	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	111	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
Peptide Transporter System	112	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	113	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	114	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	115	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	116	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	117	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	118	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	119	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	120	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	121	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	122	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	123	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	124	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	125	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	126	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	127	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	128	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	129	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	130	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
Other / Unknown	131	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	132	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	133	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	134	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	135	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	136	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	137	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	138	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	139	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	140	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	141	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	142	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	143	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	144	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	145	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	146	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	147	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	148	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	149	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline
	150	SERI	Depression / Anxiety	Desipramine	10-150 mg/day	GlaxoSmithKline

Examples

Example 1: Measuring pKi values of test compounds

5 In Table 1, the pKi values of test compounds are given for each of the dopamine receptors, 5HT receptors, adrenergic receptors and the histamine1 receptor. The affinity of test compounds for the respective receptors has been performed according to conventional procedures known in the art.

An indication "0" means that no affinity has been measured between the test compound and the receptor.

10 The columns displaying the pKi values for the D4 and the 5-HT2A receptor are filled with dark grey. pKi values between 8 and 9 and higher than 9 are represented by light grey boxes.

15 Example 2: Foregoing pipamperon-citalopram treatment in major depressive disorder: a placebo and active controlled period finding clinical trial

Table 2 represents the set-up of a clinical trial comprising for treatment groups:

Group Plc – Active / Day 0 represents the group receiving 10 mg citalopram, twice a day, starting the first day (Day 0) of active treatment in the clinical trial. This administration regime is also indicated as the mono therapy.

20 Group Pip - Active / Day 0 represents the group receiving a combination of 4 mg pipamperon and 10 mg citalopram, twice a day, starting the first day (Day 0) of active treatment in the clinical trial. This administration regime is also indicated as the non-foregoing combo therapy.

25 Group Pip - Active / Day 4 represents the group receiving 4 mg pipamperon, twice a day, starting the first day (Day 0) of active treatment in the clinical trial, followed by a combination of 4 mg pipamperon and 10 mg citalopram, twice a day, starting the fifth (Day 4) day of active treatment in the clinical trial. This administration regime is also indicated as the foregoing therapy with combination therapy starting after 4 days of active treatment.

30 Group Pip - Active / Day 7 represents the group receiving 4 mg pipamperon, twice a day, starting the first day (Day 0) of active treatment in the clinical trial, followed by a combination of 4 mg pipamperon and 10 mg citalopram, twice a day, starting the eight (Day 7) day of active treatment in the clinical trial. This administration regime is also

indicated as the foregoing therapy with combination therapy starting after 7 days of active treatment.

All subjects also undergo a placebo (PLC) run-in therapy, administered during a period of about 7 days before the active treatment starts.

- 5 During daily (D), weekly (W) or monthly (M) visits, several parameters are measured. Under NECT is to be understood: Neuronal E-clinical Trial = Vesalius Expert development for this trial which includes the bottom-up measurement of:

- In- and exclusion-criteria
- Functional status evaluation
- 10 - Medical history
- (Pre-)treatment signs & symptoms
- DSM-IV rules for diagnosis & efficacy
- HDRS-28 (Hamilton Depression Rating Scale - 28 items)
- Medical resource utilisation
- 15 - Pre-trial & Concomittant medication
- Drug administration
- (Serious) Adverse events
- Admission to the acute and extension phase of treatment
- Right flow of the trial
- 20

Example 3: combo pipamperon-citalopram: therapeutic use in Major Depression

25 Purpose

- Pipamperon (1'-[3-(p-Fluorobenzoyl)propyl]-[1,4'-bipiperidine]-4'-carboxamide), the active ingredient of Dipiperon (Janssen-Cilag B.V), administered to patients in a dose ranging between 8 and 12 mg is claimed via its specific pharmacological properties to be a booster of the antidepressant effect of the selective serotonin re-uptake inhibitor
- 30 citalopram. Preferably, pipamperon is administered daily at least 4-5 days before administering said antidepressant. The mechanism of boosting of pipamperon has to deal with (i) the selective affinity for the dopamine-4 (D4) receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other dopamine receptors, and (ii) the selective affinity for the 5-HT2A receptor with a pKi
- 35 value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards other 5HT receptors. This semi-naturalistic open label study investigated the efficacy and tolerability of the combo pipamperon - citalopram in the treatment of patients with major depression.

Details

- Design:** Semi-naturalistic i.e. inclusion of every 'natural' patient in an outpatient practice but without concomitant use of mood enhancing drugs, open label
- Control:** No
- Phase:** Phase IIa – preliminary Proof of Concept
- Location:** Belgium – Research Centre ANIMA, Aiken
- End Points :** Assessment scale scores, Hamilton Depression Rating Scale 17 items, Reduction, Response, Remission
- Medication:** Exclusion of mood stabilisers, antipsychotics (typical and atypical) and other antidepressants

Subjects

Type	No.	Sex	Age
Patients	23	10 male & 13 female	23-80 (mean 47) years

- 5 **Characteristics:** patients had a major depressive disorder according to DSM-IV criteria, with or without a chronic course and a treatment refractory state towards another SSRI then citalopram.

Treatments

- 10 **PIP-CIT¹ add-on:** citalopram from day minus 60-20 – pipamperon from DAY 0

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperon ¹	+ Pip.: 8-12 mg/day – Cit.: 20-	PO	bid	8 weeks
Citalopram ¹	40 mg/day			

1. Pipamperon (Pip) and citalopram (Cit) dosage was adjusted according to clinical response.

15

- PIP-CIT¹ fore-going 1-5:** pipamperon from day 0 – cital from day 1-5

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperon ¹	+ Pip.: 8-12 mg/day – Cit.: 20-	PO	bid	8 weeks
Citalopram ¹	40 mg/day			

- 20 1. Pipamperon (Pip) and citalopram (Cit) dosage was adjusted according to clinical response.

PIP-CIT¹ fore-going 6-8: pipamperon from day 0 – citalopram from day 6-8

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperon ¹ Citalopram ¹	+ Pip.: 8-12 mg/day – Cit.: 20-40 mg/day	PO	bid	8 weeks

- 5 1. Pipamperon (Pip) and citalopram (Cit) dosage was adjusted according to clinical response.

Results

	PIP-CIT add-on	PIP-CIT foregoing	
	After 20-60 DAYS (mean 33) (n = 5)	1-5 DAYS (mean 4) (n = 15)	6-8 DAYS (mean 7) (n = 3)
Mean Used Medication			
Pipamperone	9mg/day	10mg/day	11mg/day
Citalopram	30mg/day	26mg/day	30mg/day
Depression scale scores			
HDRS 17-item total score			
baseline	29	23	28
endpoint (week 8)	4	5	11
diminishment at week 8	-25 (+8/-9)	-18 (+8/-8)	-17 (+17/-17)
% reduction at week 8	86 (+14/-12)	80 (+20/-30)	61 (+39/-61)
response ¹ at week 8	5 (100%)	15 (100%)	2 (67%)
remission ² at week 8	4 (80%)	10 (67%)	1 (33%)

- 10 1. Response = $\geq 50\%$ reduction in HDRS 17-item score;
2. Remission = HDRS 17-item score < 8

Notably, the results obtained are highly significant since the variability in every group is distributed evenly around the mean.

15

Add-on PIP-CIT

Figure 1 schematically depicts the "add-on" treatment with pipamperon 8-12 (mean 9) mg (bid) after treatment with citalopram 10-20 (mean 30) mg (bid) during 20-60 (mean 33)

days (PIPCIT ADD-ON) with HDRS-17. Totalscore is 29 at baseline in MDD in comparison with the standard efficacy of antidepressants in clinical trials according to Khan *et al.* (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 2 schematically depicts the HDRS-17 change from baseline in the combo pipamperon as "add-on" to citalopram vs SNRI (duloxetine) in Major Depression. Treatment with pipamperon 8-12 (mean 9 mg/day) during 20-60 (mean 33) days after treatment with SSRI (n=5). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein *et al.*, (Clin. Psychiatry, in press).

Figure 3 schematically depicts the remission rates (HDRS-17 ≤ 7) with the combo pipamperon as "add-on" to citalopram vs SNRI (venlafaxine) vs SSRIs vs placebo in Major Depression. Treatment with pipamperon 8-12 (mean 9 mg/day) during 20-60 (mean 33) days after treatment with SSRI (n=5). Treatment with the SNRI venlafaxine is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241). Treatment with SSRIs is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241). Treatment with placebo is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241).

Fore-going 1-5 PIP-CIT

Figure 4 schematically depicts the "fore-going" treatment during 1-5 (mean 4) days with pipamperon 8-12 (mean 10) mg (bid), followed with the combination treatment of pipamperon and citalopram 20-50 (mean 26) mg/day (bid) (PIPCIT FG 1-5) in MDD (HDRS-17 at BL = 23) in comparison with the standard efficacy of antidepressants in clinical trials according to Khan *et al.* (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 5 schematically depicts the HDRS-17 change from baseline in the combo pipamperon-citalopram treatment with a "fore-going" treatment of 4 days with pipamperon (10 mg/day) vs SNRI (duloxetine) in Major Depression. Treatment with the combo pipamperon-citalopram with pipamperon 8-12 (mean 10 mg/day) (bid) 1-5 (mean 4) days before treatment with SSRI (n=15). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein *et al.*, (Clin. Psychiatry, in press).

Figure 6 schematically depicts the remission rates (HDRS-17 ≤ 7) with the combo pipamperon with a "fore-going" treatment of 4 days with pipamperon (10 mg/day) vs SNRI (venlafaxine) in Major Depression. Treatment with the combo pipamperon-citalopram was with pipamperon 8-12 (mean 10 mg/day) during 1-5 (mean 4) days before treatment with the SSRI (n=5). Treatment with the SNRI venlafaxine is according to a meta-analysis of Thase et al. (Br. J. Psychiatry (2001) 178:234-241). Treatment with SSRIs is according to a meta-analysis of Thase et al. (Br. J. Psychiatry (2001) 178:234-241). Treatment with placebo is according to a meta-analysis of Thase et al. (Br. J. Psychiatry (2001) 178:234-241).

Fore-going 6-8 PIP-CIT

Figure 7 schematically depicts the "fore-going" treatment during 6-8 (mean 7) days with pipamperon 8-12 (mean 11) mg/day (bid), followed with the combination treatment of pipamperon and citalopram 20-40 (mean 30) mg/day (bid) (PIPCIT FG 6-8) in MDD (HDRS-17 at BL = 28) in comparison with the standard efficacy of antidepressants in clinical trials according to Khan et al. (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 8 schematically depicts the HDRS-17 change from baseline in the combo pipamperon-citalopram treatment with a "fore-going" treatment of 7 days with pipamperon (11 mg/day) vs SNRI (duloxetine) in Major Depression. Treatment with the combo pipamperon-citalopram with pipamperon 8-12 (mean 11 mg/day) (bid) 6-8 (mean 7) days before treatment with SSRI (n=3). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein et al., (Clin. Psychiatry, in press).

Comparison "add-on" vs "fore-going"

Figure 9 schematically depicts a comparison between "fore-going" and "add-on" treatments with pipamperon (8-12 mg/day; bid) and citalopram (20-40 mg/day; bid) in MDD in comparison with the standard efficacy of antidepressants in clinical trials according to Khan et al. (2000), in "Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials" (Arch. of General Psychiatry, Vol. 57, April 2000).

Figure 10 schematically depicts a comparison between "fore-going" and "add-on" treatments. In particular, the HDRS-17 change from baseline between "fore-going" and "add-on" treatment with pipamperon (8-12 mg/day; bid) and citalopram (20-40 mg/day; bid) in comparison with the SNRI duloxetine in Major Depression is depicted. Treatment with the combo pipamperon as "add-on" to citalopram, with pipamperon 8-12 mg/day (mean 9 mg/day) 20-60 (mean 33) days after treatment with the SSRI (n=5). Treatment with the combo pipamperon-citalopram, with pipamperon 8-12 mg/day (mean 11 mg/day; bid) 6-8 days (mean 7 days) before treatment with the SSRI (n = 15). Treatment with the combo pipamperon-citalopram, with pipamperon 8-12 mg/day (mean 10 mg/day; bid) 1-5 days (mean 4 days) before treatment with the SSRI (n = 15). The SNRI (duloxetine) treatment was 40-120 mg/day (n = 152) according to Goldstein *et al.*, (Clin. Psychiatry, in press).

Figure 11 schematically depicts the remission rates (HDRS-17 ≤ 7) in a comparison between "fore-going" and "add-on" treatment with pipamperon (8-12 mg/day; bid) and citalopram (20-40 mg/day; bid) in comparison with the SNRI venlafaxine in Major Depression. Treatment with the combo pipamperon-citalopram was with pipamperon 8-12 (mean 10 mg/day) during 1-5 (mean 4) days before treatment with the SSRI (n = 15). Treatment with the SNRI venlafaxine is according to a meta-analysis of Thase *et al.* (Br. J. Psychiatry (2001) 178:234-241). Treatment with pipamperon as "add-on" to citalopram, with pipamperon 8-12 (mean 9 mg/day) during 20-60 (mean 33) days after treatment with SSRI (n = 5).

The intention-to-treat / last-observation-carried-forward analysis showed a high therapeutic efficacy according HDRS 17-item in all the treatment groups. This was especially true for the 'add-on' group probably caused by the longer treatment with an active antidepressant (+33 days). The huge therapeutic effect observed in the 'PIP-CIT 1-5' group present for at a mean dosage of pipamperon of 10 mg per day and administered the first four days of treatment without an active antidepressant, indicates the boosting effect of pipamperon on the SSRI citalopram at an extremely and thus unconventional low dose. Only 1 patient discontinued treatment due to a lost of follow-up.

201

Adverse Events

Side effects (patients)	PIP-CIT add-on After 20-60 DAYS (mean 33) (n = 5)	PIP-CIT foregoing 1-5 DAYS (mean 4) (n = 15)	6-8 DAYS (mean 7) (n = 3)
Discontinued treatment due to adverse events	0	0	0
By system:			
body as a whole	0	0	0
central and peripheral nervous system	1(20%)	4(26.6%)	0
gastrointestinal	1(20%)	5(33%)	2(66.6%)
musculoskeletal	1(20%)	3(20%)	0
psychiatric	0	0	0
respiratory	0	1(6.6%)	0
skin and appendages	1(20%)	2(13.3%)	1(33.3%)
vascular	0	1(6.6%)	0
urinary	0	1(6.6%)	0

Laboratory parameters, ECG, bodyweight and vital signs were not measured since this was a naturalistic study.

5

Assessment**Outcome**

Efficacy: the 4-day fore-going combo pipamperon 8-12mg/d - citalopram 20-40mg/day is comparable to the add-on combo pipamperon-citalopram.

10 **Efficacy:** the 4-day fore-going combo pipamperon 8-12mg/d - citalopram 20-40mg/day is larger than the 7-day fore-going combo pipamperon 8-12mg/d - citalopram 20-40mg/day.

Efficacy: the combo pipamperon 8-12mg/d - citalopram 20-40mg/day is larger than the in the art known antidepressants SSRIs.

15

Tolerability

Tolerability: the 4-day fore-going treatment is comparable to the 7-day fore-going combo is comparable to add-on combo pipamperon-citalopram.

Tolerability: no discontinued treatment due to adverse events.

5

Study Messages

The boosting effect of pipamperon at an extremely unconventional low dose on a SSRI is indicated since the efficacy of the 'add-on' and '4-day fore-going' combo 'pipamperon 8 -12 mg/d - citalopram 20 - 40 mg/day' is in this study as twice higher as known in the art in the treatment of patients with major depression.

10

The combo pipamperon-citalopram is generally well tolerated in patients with depression i.e. at least no specific added adverse events were occurring by adding pipamperon at the doses used in the study.

15

Example 4: combo pipamperon-citalopram: therapeutic use in Obsessive-Compulsive Disorder (OCD).**Purpose**

- 5 Pipamperon (1'-[3-(p-Fluorobenzoyl)propyl]-[1,4'-bipiperidine]-4'-carboxamide), the active ingredient of Dipiperon (Janssen-Cilag B.V), administered to a patient in a dose ranging between 8 and 12 mg is claimed via its specific pharmacological properties to be a booster of the effect of the selective serotonin re-uptake inhibitor citalopram towards OCD. Preferably, pipamperon is administered daily at least 4-5 days before
- 10 administering said antidepressant. The mechanism of boosting of pipamperon has to deal with (i) the selective affinity for the dopamine-4 (D4) receptor with a pKi value equal to or higher than 8 towards the D4 receptor and less than 8 towards other Dopamine receptors, and (ii) the selective affinity for the 5-HT2A receptor with a pKi value equal to or higher than 8 towards the 5-HT2A receptor and less than 8 towards
- 15 other 5HT receptors. This semi-naturalistic open label study investigated the efficacy and tolerability of the combo pipamperon - citalopram in the treatment of patients with OCD.

Details

- Design: Semi-naturalistic i.e. Inclusion of every 'natural' patient in an outpatient practice but without concomitant use of mood enhancing drugs, open label
- Control: No
- Phase: Phase IIa – preliminary Proof of Concept
- Location: Belgium – Research Centre ANIMA, Alken
- End Points : Assessment scale scores, Yale-Brown Obsessive-Compulsive Scale, Reduction, Remission
- Medication: Exclusion of mood stabilisers, antipsychotics (typical and atypical) and other antidepressants

20

Subjects

Type	No.	Sex	Age
Patients	7	1 male & 7 female	20-63 (mean 33) years

Characteristics: patients had an obsessive-compulsive disorder according to DSM-IV criteria, with or without a chronic course and a treatment refractory state towards another SSRI then citalopram.

5 Treatments

PIP-CIT¹ ADD-ON: citalopram from DAY minus 730-60 – pipamperon from DAY 0

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperone ¹	+ Pip.: 8-16 mg/day – Cit.: 30-80	PO	bid	12
Citalopram ¹	mg/day			weeks

- 10 1. Pipamperone (Pip) and Citalopram (Cit) dosage was adjusted according to clinical response.

PIP-CIT¹ FORE-GOING 4-6: pipamperon from DAY 0 – citalopram from DAY 4-6

Drug/Treatment	Dose	Route	Frequency	Duration
Pipamperone ¹	+ Pip.: 8-16 mg/day – Cit.: 30-80	PO	bid	12
Citalopram ¹	mg/day			weeks

15

1. Pipamperone (Pip) and Citalopram (Cit) dosage was adjusted according to clinical response.

Results

PIP-CIT add-on	
after 730-60 DAYS (mean 241) (n = 6)	
with mean Cit. 54mg/d and Pip. 11mg/d	
PIP-CIT foregoing	
4-6 DAYS (mean 5) (n = 2)	
with mean Cit. 60mg/d and Pip. 10mg/d	
Y-BOCS score	
<u>Baseline</u>	
Total	31
Obsessions	18
Compulsions	13
Endpoint (week 12)	
<u>Total</u>	<u>15</u>
diminishment	-16 (+16/-11)
% reduction	53
<u>Obsessions</u>	
total	8
diminishment	-10 (+9/-7)
% reduction	57
<u>Compulsions</u>	
total	7
diminishment	-6 (+7/-6)
% reduction	45
<u>% Remission</u>	
YBOCS score ≤8	29
BOCS score ≤16	57

Notably, the results obtained are highly significant since the variability in every group is distributed evenly around the mean.

Figure 12 schematically depicts the Y-BOCS total score: "fore-going" and "add-on" treatment with pipamperon (8-15 mg/day; bld) and citalopram (30-80 mg/day; bid) in comparison with the SSRI fluvoxamine in OCD. Treatment with the combo pipamperon-

citalopram (n = 7). Treatment with fluvoxamine (controlled release) mean 271 mg/day (n = 253) is according to Hollander et al. (2003).

Figure 13 schematically depicts the Y-BOCS obsession score: "fore-going" and "add-on" treatment with pipamperon (8-15 mg/day; bid) and citalopram (30-80 mg/day; bid) in comparison with the SSRI fluvoxamine in OCD. Treatment with the combo pipamperon-citalopram (n = 7). Treatment with fluvoxamine (controlled release) mean 271 mg/day (n = 253) is according to Hollander et al. (2003).

Figure 14 schematically depicts the Y-BOCS compulsion score: "fore-going" and "add-on" treatment with pipamperon (8-16 mg/day; bid) and citalopram (30-80 mg/day; bid) in comparison with the SSRI fluvoxamine in OCD. Treatment with the combo pipamperon-citalopram (n = 7). Treatment with fluvoxamine (controlled release) mean 271 mg/day (n = 253) is according to Hollander et al. (2003):

The intention-to-treat / last-observation-carried-forward analysis showed a high therapeutic efficacy according Y-BOCS total score, obsession and compulsion scores. This indicates the boosting effect of pipamperon on the SSRI citalopram at an extremely and thus unconventional low dose. No patient discontinued treatment.

Assessment

Efficacy: the combo pipamperone 8-16mg/d - citalopram 30-80mg/day > the in the art known compounds effective towards OCD (Hollander E, Koran LM, Goodman WK, Greist JH, Ninan PT, et al. A double-blind, placebo-controlled study of the efficacy and safety of controlled-release fluvoxamine in patients with obsessive-compulsive disorder. Journal of Clinical Psychiatry 64: 640-647, Jun 2003 Mount Sinai School of Medicine, New York, New York, USA; Solvay Pharmaceuticals Inc., Marietta, Georgia, USA).

Study Messages

The boosting effect of pipamperon at an extremely unconventional low dose on a SSRI is indicated since the efficacy of the 'add-on' and 'fore-going' combo 'pipamperon 8-15 mg/d - citalopram 30-80 mg/day' is in this study as twice higher as known in the art in the treatment of patients with obsessive-compulsive disorder.

Example 5: combo pipamperon-citalopram: therapeutic use in Panic Disorder.**Purpose**

- 5 Preliminary examination of a "fore-going" and "add-on" treatment with pipamperon and citalopram in comparison with the SSRI in Panic Disorder.

Results

- 10 The results are indicated in Figure 15. Figure 15 schematically depicts the CGI-severity score: "fore-going" and "add-on" treatment with pipamperon (8 mg/day; bid) and citalopram (20-40 mg/day; bid) in comparison with the SSRI in Panic Disorder. Treatment with the combo pipamperon-citalopram ($n = 3$). Treatment with paroxetine is according to the Journal of Clinical Psychiatry (2004) 65: 405-413. Treatment with Sertraline is according to the Journal of Clinical Psychiatry (2004) 65: 405-413.

15 Conclusion

- 20 Notably, although a small test group has been used ($n = 3$), the distribution around the mean is good. It will further be apparent from Figure 15 that the effect of the combo treatment of pipamperon and citalopram is twice as high as the standard treatments with paroxetine or sertraline.

Example 6: POC process for mayor depressive disorder

Concept: Combo of the high selective 5-HT_{2A/D4} antagonist pipamperon with:

- 25 - a compound active towards the Amino Acid Transmitter, Peptidergic Transmitter, Adenosine Transmitter, Endocrine and/ or Enzymatic System;
- a fore-going admission during 4 days of pipamperon;
- a dose of pipamperon of 12 mg/day

30 Objectives: Demonstrating that this combo therapy has:

- 35 - the potency of being a treatment standard for depression by having an added value of reducing the total score of the Hamilton Depression Rating Scale – 17 items (HDRS-17) after 8 weeks of therapy with a least 20% more than reached with the conventional known antidepressants, i.e. 60% versus 40%. This stands for an added medium demission of 5 points on the total score of the HDRS-17 and by this will be very highly

significant since the mean difference in all recent clinical trials between placebo and active treatment is 2.5;

- a more sustained therapeutic effect than the conventional mono therapy by preventing significant more relapses during 48 weeks following the acute treatment; and/or
- 5 - a complete neutral safety profile, e.g. there are no more adverse events in the combo therapy than in mono admission of the in the combo used antidepressant compound.

Process: the following different steps were implemented to reach out for these objectives (see also Tables 3 and 4):

- 10 (1) an naturalistic open label study ($n \geq 20$) on a depressive population with a normal variability of medical and psychiatric history, course of depression, earlier and concomitant therapy admitting the golden standard antidepressant citalopram 20-40 mg/day and a dose of 8-12 mg/day of pipamperon in a foregoing, simultaneous or add-on use.

- 15 (2) a 16 weeks placebo controlled randomised four armed study of each 36 patients with a mayor depressive disorder admitting:

- from day 0: placebo or pipamperon (PIP) 10 mg/day or an active antidepressant compound or the combination of the last two;
 - from day 4: placebo or pipamperon 10 mg/day combined with an active antidepressant compound or an active antidepressant compound without pipamperon.
- 20

By including rigorous control groups (placebo and active comparator; see Tables 3 and 4) this clinical trial is evaluated as a proof of concept of the added value of the combo and the foregoing treatment method since the inclusion/exclusion of:

- 25
- a negative trial, i.e. no significant difference between the placebo and active treatment with the comparator;
 - a failed trial, i.e. no significant difference between the active and the studied treatment i.e. the combo.

- 30 (3) an active controlled randomised relapse prevention study following the POC trial during another 36 weeks with three arms of each 36 patients which is formed by:

- continuation of the active mono therapy;
- randomising the patients with a combo therapy in a group with an active mono therapy and with a continuation of the combo treatment.

Claims

1. Use of pipamperon for the preparation of a medicament for treating a disease or disorder with an underlying dysregulation of the emotional functionality, wherein said pipamperon is administered to a patient in a dose ranging between 5 and 15 mg of the active ingredient, and wherein a second compound is administered simultaneously with, separate from or sequential to said pipamperon to augment the therapeutic effect of said second compound.
2. Use according to claim 1, wherein said pipamperon is administered daily at least one day before administering said second compound.
3. Use according to any of claims 1 to 2, wherein said second compound affects the monoaminergic transmitter system.
4. Use according to claim 3, wherein said second compound is selected from the group comprising: 5-HT reuptake enhancer (1), 5-HT1 autoreceptor agonist (2), 5HT1A receptor agonist (3), 5-HT1A receptor antagonist (4), 5-HT1B receptor antagonist (5), 5-HT2B receptor antagonist (6), 5-HT2C receptor antagonist (7), 5-HT3 receptor antagonist (8), 5-HT6 receptor antagonist (9), adrenergic transmitter releaser (12), α 1 adrenoreceptor antagonist (13), α 2 adrenoreceptor antagonist (14), β 3 adrenoreceptor agonist (19), cannaboid receptor antagonist (21), D1 receptor agonist (27), D2 receptor antagonist (28), D3 receptor antagonist (29), DA uptake inhibitor (30), dopamine receptor agonist (31), H3 receptor antagonist (42), compounds which increase brain concentrations of 5-HT (44), levodopa (48), MAO reuptake inhibitor (50), MAO-A & MAO-B reuptake inhibitor (51), MAO-B inhibitor (52), MAO-B re-uptake inhibitor (53), NARI (60), NaSSA (61), NDRI (62), RIMA (82), SDA (84), SDRI (85), Second messenger beta agonist (86), SNDRI (90), SNRI (91) and SSRI (92).
5. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial

behaviour, bereavement, occupational problem and problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT (serotonin) reuptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT (serotonin) reuptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10 6. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases or disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₁ autoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₁ autoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25 7. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-induced persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic

disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder, other cognitive disorders and Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A) receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin) 1A receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

8. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

9. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1B} receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1B} receptor

antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10. Use of pipamperon or a pharmaceutically acceptable salt thereof for the
5 preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse
10 control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2B} receptor antagonist compound to augment the therapeutic effect or to provide a
15 faster onset of the therapeutic effect of said 5-HT_{2B} receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

11. Use of pipamperon or a pharmaceutically acceptable salt thereof for the
20 preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender disorders, sleep disorders, adjustment disorders, impulse
25 control disorders, substance-related disorder, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{2C} receptor antagonist compound to augment the therapeutic effect or to provide a
30 faster onset of the therapeutic effect of said 5-HT_{2C} receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

12. Use of pipamperon or a pharmaceutically acceptable salt thereof for the
35 preparation of a medicament for treating the underlying emotion dysregulation of substance-related disorders, characterized in that pipamperon or said pharmaceutically

acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₃ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₃ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

13. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive disorder selected from the group of diseases and disorders consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₆ receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₆ receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

14. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an adrenergic transmitter releaser compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said adrenergic transmitter releaser compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a alpha 1 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 1 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

16. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a alpha 2 adrenoreceptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said alpha 2 adrenoreceptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

17. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders,

- impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of
- 5 a beta 3 adrenoreceptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said beta 3 adrenoreceptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
- 10 18. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders,
- 15 adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, pain disorders and delirium, characterized in that pipamperon or said
- 20 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a cannaboid receptor 1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said cannaboid receptor 1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
- 25 ingredient.
19. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders and Parkinson disease, characterized in that pipamperon or
- 30 said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D1 receptor receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D1 receptor receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
- 35 ingredient.

20. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, pain disorders and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D2 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D2 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

21. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorders and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a D3 receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said D3 receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

22. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of substance related disorders and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with,

5 separate from or prior to the administration of a DA (dopamine) uptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said DA (dopamine) uptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10 23. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders and problems related to abuse or neglect, pain disorder and Parkinson disease, characterized in that pipamperon or said
15 pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a dopamine receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said dopamine receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active
20 ingredient.

24. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a cognitive mental disease or disorder selected from the group of diseases and disorders
25 consisting of Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment
30 disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a histamine H3receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said histamine H3receptor antagonist compound, further characterized in that pipamperon is to
35 be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases brain concentrations of 5-HT (serotonin) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases brain concentrations of 5-HT (serotonin), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

26. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a levodopa/decarboxylase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said levodopa/decarboxylase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

27. A pharmaceutical composition comprising (a) pipamperon, and (b) a levodopa/decarboxylase inhibitor compound, preferably is (eti)levodopa/carbidopa, or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof in combination with entacapone, which is an inhibitor of catechol-O-methyltransferase (COMT), or a pro-drug or an active metabolite thereof, or a pharmaceutically acceptable salt thereof, as a combined preparation for simultaneous, separate or sequential use for treating the underlying emotion dysregulation of Parkinson Disease.

28. A pharmaceutical composition according to claim 27, wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein

said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa, preferably provided in a unitary dose of between 100 mg and 10 mg of the active ingredient.

29. A pharmaceutical composition according to claim 27 or 28, wherein pipamperon is provided in a unitary dose of between 5 and 15 mg of the active ingredient and wherein said levodopa/decarboxylase inhibitor compound is levodopa/carbidopa or etilevodopa/carbidopa in combination with entacapone, of which the latter is preferably provided in a unitary dose of between 500 mg and 100 mg of the active ingredient.

30. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of non-cognitive mental disease or disorder which are substance related disorders and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase (MAO) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase (MAO) reuptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

31. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase A (MAO-A) and a monoamine oxidase B (MAO-B) reuptake inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

32. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, problems related to abuse or neglect, pain disorder and Parkinson disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

33. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Parkinson Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a monoamine oxidase B (MAO-B) reuptake inhibitor to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said monoamine oxidase B (MAO-B) reuptake inhibitor, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

34. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorder, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline re-uptake inhibitor (NARI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline re-uptake

inhibitor (NARI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a noradrenergic/specific serotonergic antidepressant (NaSSA) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said noradrenergic/specific serotonergic antidepressant (NaSSA) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
36. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, adjustment disorders, attention-deficit disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
37. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders

consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, antisocial behaviour, bereavement, occupational
5 problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which is a reversible inhibitor of mono-amine oxydase A (RIMA) to augment the therapeutic effect or
10 to provide a faster onset of the therapeutic effect of said compound which is a reversible inhibitor of mono-amine oxydase A (RIMA), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

38. Use of pipamperon or a pharmaceutically acceptable salt thereof for the
15 preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development
20 disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem, problems related to abuse or neglect, pain disorder and delirium, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate
25 from or prior to the administration of a serotonin/dopamine antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said serotonin/dopamine antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30

39. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome,
35 somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-

related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect, pain disorders, delirium, Alzheimer Disease, substance-related persisting dementia, vascular dementia, dementia due to HIV disease, dementia due to head trauma, dementia due to Parkinson Disease, dementia due to Huntington Disease, dementia due to Pick Disease, dementia due to Creutzfeldt-Jacob Disease, amnesic disorders due to a general medical condition, substance-induced persisting amnesic disorder, mild cognitive impairment disorder and other cognitive disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine re-uptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine re-uptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

40. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a second messenger beta agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said second messenger beta agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

41. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-

deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

42. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, attention-deficit disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and nor-adrenaline re-uptake inhibitor (SNRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

43. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, antisocial behaviour, bereavement, occupational problem, problems related to abuse or neglect and pain disorders, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin

225

re-uptake inhibitor (SSRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin-re-uptake inhibitor (SSRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

5

44. Use according to claim 1 or 2, wherein said disease or disorder is Alzheimer disease.

10

45. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT_{1A} (serotonin 1A receptor) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT_{1A} (serotonin 1A receptor) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

15

20

46. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a 5-HT₆ antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said 5-HT₆ antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

25

30

47. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an acetylcholinesterase inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said acetylcholinesterase inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

35

226

48. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an AMPA receptor mediator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said AMPA receptor mediator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
49. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an amyloid aggregation-inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said amyloid aggregation-inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
50. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a calcium channel modulator compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said calcium channel modulator compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.
51. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a choline uptake enhancer compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said choline uptake enhancer compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

52. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound that activates ERK (extracellular signal-related kinase) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound that activates ERK (extracellular signal-related kinase), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

10

53. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GABA agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GABA agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

20

54. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a histamine H3-receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said histamine H3-receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

30

55. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which increases insulin sensitivity to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which increases insulin sensitivity, further characterized in that pipamperon is to be

35

228

administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

56. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which mimics the effect of nerve growth factor (NGF) to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which mimics the effect of nerve growth factor (NGF), further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

57. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a muscarinic receptor partial agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said muscarinic receptor partial agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

58. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective nor-adrenaline and dopamine re-uptake inhibitor (NDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

59. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable

salt thereof is administered simultaneously with, separate from nerve growth factor (NGF) gene therapy, to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nerve growth factor (NGF) gene therapy, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

60. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a nicotinic receptor agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said nicotinic receptor agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

61. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of an N-Methyl-D-aspartate (NMDA) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said N-Methyl-D-aspartate (NMDA) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

62. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a non-steroidal anti-inflammatory drug to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said a non-steroidal anti-inflammatory drug, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

63. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of

Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a phosphodiesterase-4 (PDE4) inhibitor compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said phosphodiesterase-4 (PDE4) inhibitor compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

64. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

65. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a compound which protects dopaminergic and cholinergic neurons to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said compound which protects dopaminergic and cholinergic neurons, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

66. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

67. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin and dopamine reuptake inhibitor (SDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin and dopamine reuptake inhibitor (SDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

68. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of Alzheimer Disease, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said selective serotonin, nor-adrenaline and dopamine re-uptake inhibitor (SNDRI) compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

69. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance related disorders, personality disorders, bereavement, occupational problem and ~~patterns~~ related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a CRF1 antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said CRF1 antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

70. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a GR antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said GR antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

71. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a melatonin receptor (MT) agonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said melatonin receptor (MT) agonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

72. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep

disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurotensin receptor antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurotensin receptor antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

73. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 2 receptor (NK2) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 2 receptor (NK2) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

74. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, psychotic disorders, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sleep disorders, adjustment disorders, impulse control disorders, pervasive development disorders, disruptive behaviour disorders, substance-related disorders, personality disorders, psychological factors affecting medical conditions, malingering, antisocial behaviour, bereavement, occupational problem, identity problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable

234

salt thereof is administered simultaneously with, separate from or prior to the administration of a neurokinin 3 receptor (NK3) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said neurokinin 3 receptor (NK3) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

75. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, adjustment disorders, impulse control disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a peptidic compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said peptidic compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

76. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a substance P receptor (NK1) antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said substance P receptor (NK1) antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

235

77. Use of pipamperon or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treating the underlying emotion dysregulation of a non-cognitive mental disease or disorder selected from the group of diseases and disorders consisting of mood disorders, anxiety disorders, eating disorders, premenstrual syndrome, somatoform disorders (excluding pain disorders), factitious disorders, dissociative disorders, sexual and gender identity disorders, sleep disorders, adjustment disorders, impulse control disorders, substance-related disorders, personality disorders, bereavement, occupational problem and problems related to abuse or neglect, characterized in that pipamperon or said pharmaceutically acceptable salt thereof is administered simultaneously with, separate from or prior to the administration of a tachykinin antagonist compound to augment the therapeutic effect or to provide a faster onset of the therapeutic effect of said tachykinin antagonist compound, further characterized in that pipamperon is to be administered to a patient in a daily dose ranging between 5 and 15 mg of the active ingredient.

78. Use according to any of claims 1 to 4 or 7, wherein said second compound is gepirone.

79. A method for preparing a compound having a selective D4 and 5-HT2A antagonist, reverse agonist or partial agonist activity comprising the following steps: (a) measuring the selective affinity of a test compound to the D4 receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the D4 receptor in respect to all the other D receptors, and measuring the selective efficacy of the selected compound to the D4 receptor and selecting a compounds which is a selective antagonist, inverse agonist or partial agonist of the D4 receptor; (b) measuring the selective affinity of a test compound to the 5-HT2A receptor and selecting a compound that has a pKi value equal to or greater than 8 towards the 5-HT2A receptor in respect to all the other 5HT receptors, and measuring the selective efficacy of the selected compound to the 5-HT2A receptor and selecting a compounds which is a selective antagonist, inverse agonist or partial agonist of the 5-HT2A receptor; (c) identifying a compound which is selected in (a) and (b), (d) preparing the compound identified in (c).

80. Compound prepared by the method of claim 79.

81. Use according to any of claims 1, 2 or 7, wherein said second compound is chosen from the group consisting of fluvoxamine controlled release, phenserine tartrate,

236

atomoxetine hydrochloride, bupropion (controlled-release formulation), ropinirole HCL (controlled-release formulation), INN 00835, galantamine (extended release formulation), paliperidone, tomoxetine, aprepitant, rivastigmine tartrate, ORG 34517/34850, sunepitron, sumanirole, milnacipran, idazoxan, xaliproden, SR 58611, befloxatone, litoxetine, 5 tianeptine, agomelatine, SPD 503, flesinoxan, bifeprunox, ramelteon, etilevodopa, rasagiline (TVP-1012) and desvenlafaxine.

82. Use according to any of claims 1, 2 or 7, wherein said second compound is chosen from the group consisting of galantamine (extended release formulation), 10 R121919, risperidone, paliperidone and R228060 (YKP-10A).

Abstract


The present invention relates to the use of compounds and compositions of compounds having D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic activity for the treatment of the underlying dysregulation of the emotional functionality of mental disorders (i.e. affect instability – hypersensitivity – hyperaesthesia – dissociative phenomena - etc).

5 The invention also relates to methods comprising administering to a patient diagnosed as having a neuropsychiatric disorder a pharmaceutical composition containing (i) compounds having D4 antagonistic, partial agonistic or inverse agonistic activity and (ii) compounds having 5-HT2A antagonistic, partial agonistic or inverse agonistic, and (iii) any
10 known medicinal compound and compositions of said compounds. The combined D4 and 5-HT2A antagonistic, partial agonistic or inverse agonistic effects may reside within the same chemical or biological compound or in two different chemical and/or biological compounds.

1/15

Add-On Treatment with Pipamperon 8-12 (mean 9) mg (bid) after Treatment with Citalopram 10-20 (mean 30) mg (bid) during 20-60 (mean 33) days (PICIT ADD-ON) with HDRS-17 Totalscore = 29 at Baseline in MDD in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*

HDRS-17 REDUCTION OVER 8 WEEKS IN MDD



A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, *ARCH. OF GENERAL PSYCHIATRY* / VOL 57, APR 2000

Figure 1

2/15

HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON AS ADD-ON - CITALOPRAM vs SNRI (duloxetine) in MAJOR DEPRESSION




Figure 2

3/15

REMISSION RATES (HDRS-17 ≤ 7): COMBO PIPAMPERON AS ADD-ON - CITALOPRAM vs SNRI (venlafaxine) vs SSRI's vs PLACEBO in MAJOR DEPRESSION





Figure 3

4/15

Foregoing Treatment During 1-5 (mean 4) days with Pipamperon 8-12 (mean 10) mg/day (bid) Followed With the Combination Treatment of Pipamperon and Citalopram 20-50 (mean 26) mg/day (bid) (PICIT FG 1-5) in MDD (HDRS-17 at BL = 23) in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*

HDRS-17 REDUCTION OVER 8 WEEKS IN MDD



A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCHIVES OF GENERAL PSYCHIATRY, VOL 57, APR 2000

Figure 4

5/15

HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON- CITALOPRAM WITH A FORE-GOING TREATMENT OF 4 DAYS WITH PIPAMPERON 10mg/day vs SNRI (duloxetine) in MAJOR DEPRESSION




Figure 5

6/15

REMISSION RATES (HDRS-17 ≤ 7): COMBO PIPAMPERON-CITALOPRAM WITH A FORE-GOING TREATMENT OF 4 DAYS WITH PIPAMPERON 10mg/day vs SNRI (venlafaxine) in MAJOR DEPRESSION





Figure 6

7/15

Foregoing Treatment During 6-8 (mean 7) days with Pipamperon 8-12 (mean 11) mg/day (bid) Followed With the Combination Treatment of Pipamperon and Citalopram 20-40 (mean 30) mg/day (bid) (PICIT FG 6-8) in MDD (HDRS-17 at BL = 28 in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*

HDRS-17 REDUCTION OVER 8 WEEKS IN MDD



A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCHIVES OF GENERAL PSYCHIATRY/VOL 57/APR 2000

Figure 7

8/15

HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON- CITALOPRAM WITH A FORE-GOING TREATMENT OF 7 DAYS WITH PIPAMPERON 11mg/day vs SNRI (duloxetine) in MAJOR DEPRESSION





Figure 8

Foregoing & Add-On Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in MDD in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials*

HDRS-17 REDUCTION OVER 8 WEEKS IN MDD



* A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)

Figure 3

M/15

REMISSION RATES (HDRS-17 ≤ 7): Foregoing & Add-On Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the SNRI venlafaxine in MAJOR DEPRESSION




Figure 11

12/15

Y-BOCS TOTALSCORE: Foregoing & Add-On Treatment with Pipamperon 8-16 mg/day (bid) and Citalopram 30-80 mg/day (bid) in comparison with the SSRI fluvoxamine in OCD




Figure 12

13/15

Y-BOCS OBSESSION SCORE: Foregoing & Add-On Treatment with Pipamperon 8-15 mg/day (bid) and Citalopram 30-80 mg/day (bid) in comparison with the SSRI fluvoxamine in OCD




Figure 13

14/15

Y-BOCS COMPULSION SCORE: Foregoing & Add-On Treatment with Pipamperon 8-15 mg/day (bid) and Citalopram 30-80 mg/day (bid) in comparison with the SSRI fluvoxamine in OCD





Figure 14

15/15

CGI-SEVERITY SCORE: Foregoing & Add-On Treatment with Pipamperon 8 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the SSRI in Panic Disorder



—●— Paroxetine in the treatment of panic disorder, Journal of Clinical Psychiatry 65: 405-413, No. 3, Mar 2004

—*— COMBO PIPAMPERON-CITALOPRAM (n=3)

—*— Sertraline in the treatment of panic disorder, Journal of Clinical Psychiatry 65: 405-413, No. 3, Mar 2004

Figure 15

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/BE04/000172

International filing date: 02 December 2004 (02.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 04025035.9
Filing date: 21 October 2004 (21.10.2004)

Date of receipt at the International Bureau: 14 April 2005 (14.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.